

2019-10-30

# Allele-selective lowering of mutant HTT protein by HTTLC3 linker compounds

Li, Z

<http://hdl.handle.net/10026.1/15145>

---

10.1038/s41586-019-1722-1

Nature

Springer Science and Business Media LLC

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

Allele-selective Lowering of Mutant HTT Protein by HTT-LC3 Linker Compounds

Zhaoyang Li<sup>1#</sup>, Cen Wang<sup>1#</sup>, Ziyang Wang<sup>1#</sup>, Chenggang Zhu<sup>2#</sup>, Jie Li<sup>3</sup>, Tian Sha<sup>1</sup>,  
Lixiang Ma<sup>4</sup>, Chao Gao<sup>5</sup>, Yi Yang<sup>6</sup>, Yimin Sun<sup>1</sup>, Jian Wang<sup>1</sup>, Xiaoli Sun<sup>1</sup>, Chenqi Lu<sup>1</sup>,  
Marian Difulgia<sup>7</sup>, Yanai Mei<sup>1</sup>, Chen Ding<sup>1§</sup>, Shouqing Luo<sup>6§</sup>, Yongjun Dang<sup>8</sup>, Yu Ding<sup>1\*</sup>,  
Yiyan Fei<sup>2\*</sup>, Boxun Lu<sup>1\*</sup>

<sup>1</sup> Neurology Department at Huashan Hospital, State Key Laboratory of Medical  
Neurobiology and MOE Frontiers Center for Brain Science, Institutes of Brain Science,  
School of Life Sciences, Fudan University, Shanghai, China.

<sup>2</sup> Department of Optical Science and Engineering, Shanghai Engineering Research  
Center of Ultra-Precision Optical Manufacturing, Key Laboratory of Micro and Nano  
Photonic Structures (Ministry of Education), Fudan University, Shanghai, China.

<sup>3</sup> National Facility for Protein Science in Shanghai, Large-scale Preparation System,  
Shanghai, China.

<sup>4</sup> Department of Anatomy, Histology and Embryology, Shanghai Medical College,  
Fudan University, Shanghai, China.

<sup>5</sup> Institutes of Biomedical Sciences, Fudan University, Shanghai, China.

<sup>6</sup> Peninsula Schools of Medicine and Dentistry, Institute of Translational and Stratified  
Medicine, University of Plymouth, Plymouth, UK.

<sup>7</sup> Laboratory of Cellular Neurobiology, Department of Neurology, Massachusetts  
General Hospital, Charlestown, Massachusetts, United States of America

<sup>8</sup> Key Laboratory of Metabolism and Molecular Medicine, Shanghai Medical College,  
Fudan University, Shanghai, China.

# These authors contribute equally;

\*Corresponding authors:

Boxun Lu (lead contact): [luboxun@fudan.edu.cn](mailto:luboxun@fudan.edu.cn)

Yiyan Fei: [fyy@fudan.edu.cn](mailto:fyy@fudan.edu.cn)

Yu Ding: [yuding@fudan.edu.cn](mailto:yuding@fudan.edu.cn)

§Co-senior authors

## 31 **Summary**

32 Accumulation of mutant proteins is the major cause for many diseases  
33 (proteopathies), and lowering the level of these proteins is highly desired for treatment.  
34 We hypothesized that compounds interacting with both the autophagosome protein  
35 LC3<sup>1</sup> and the disease-causing protein may target the latter for autophagic clearance.  
36 We tested this hypothesis in the context of lowering mutant HTT protein (mHTT),  
37 which contains an expanded polyglutamine (polyQ) tract and causes Huntington's  
38 disease (HD), an incurable neurodegenerative disorder<sup>2</sup>. Through  
39 small-molecule-microarray based screening, we identified four compounds interacting  
40 with both LC3 and mHTT, but not the wild-type HTT protein (wtHTT). Some of these  
41 compounds targeted mHTT to autophagosomes, reduced mHTT levels in an  
42 allele-selective manner, and rescued HD-relevant phenotypes in cells and *in vivo* in  
43 the fly and mouse HD models. We further revealed that these compounds interacted  
44 with the expanded polyQ stretch and could lower the level of mutant ATXN3, another  
45 disease-causing protein with expanded polyQ<sup>3</sup>. Our study provides candidate  
46 compounds for lowering mHTT and potentially other disease-causing proteins with  
47 polyQ expansion, demonstrating the concept of lowering disease-causing proteins by  
48 autophagosome-tethering compounds (ATTEC).

## 49    **Background**

50       An emerging approach for disease treatment is to lower the levels of  
51   disease-causing proteins, especially those with unknown activities. Biological tools  
52   such as RNAi or CRISPR may achieve this goal<sup>4-6</sup>, but their clinical delivery is  
53   challenging. Enhancing proteasomal degradation of target proteins by “PROTAC” is a  
54   promising emerging approach<sup>7</sup>, but proteasomes alone are inefficient in degrading  
55   certain large proteins or aggregates<sup>8</sup>. Another independent protein degradation  
56   pathway is macroautophagy (referred to as autophagy hereafter), which is a bulk  
57   degradation system that engulfs proteins into autophagosomes for subsequent  
58   lysosomal degradation<sup>9</sup>. Autophagy is present in all eukaryotic cells, and thus  
59   harnessing the power of autophagy to degrade certain target proteins may open new  
60   windows for drug discovery. Here we investigate this possibility in the context of  
61   lowering mHTT, which contains an expanded polyglutamine (polyQ) stretch (≥36Q)  
62   and causes HD, an incurable monogenetic neurodegenerative disorder<sup>2</sup>.

63       mHTT could be degraded by autophagy, during which protein substrates are  
64   engulfed into double-membrane autophagosomes associated with lipidated LC3<sup>1</sup>. We  
65   thus hypothesized that linker compounds interacting with both mHTT and LC3 may  
66   tether them together to enhance the recruitment of mHTT into autophagosomes,  
67   facilitating mHTT degradation. In addition, mHTT-LC3 linker compounds that do not  
68   interact with wtHTT may promote allele-selective degradation of mHTT. Since no  
69   mHTT/LC3-interacting compounds have been reported, we performed  
70   small-molecule-microarray-based screening for desired compounds, and utilized



71 wtHTT for the counter-screen to identify allele-selective candidates.

## 72 **Results**

### 73 *Identification of mHTT-LC3 linker compounds*

74 We stamped 3375 compounds (Fig. 1a) in duplicates into a microarray on  
75 isocyanate-functionalized glass slides *via* the nucleophile-isocyanate reaction, which  
76 forms covalent bonds between the compounds and the glass slides<sup>10,11</sup>. We then  
77 purified the human LC3B protein<sup>1</sup> (Extended Data Fig. 1a-b & Supplementary Table 1),  
78 a pathogenic mHTT exon1 fragment<sup>12</sup> with expanded polyQ (mHTTExon1-Q72), and a  
79 control wtHTT exon1 fragment (HTTExon1-Q25) (Extended Data Fig. 1c-d) for the  
80 screen. We fused a maltose-binding-protein (MBP) tag to both HTT exon1 proteins to  
81 increase their solubility required for later experiments.

82 To identify LC3B- and mHTT- interacting compounds, we had these purified  
83 proteins flow through the SMMs, and detected the compound-protein interaction using  
84 an optical biosensor, the scanning oblique-incidence reflectivity difference (OI-RD)  
85 microscope. OI-RD is a widely used interaction measurement technology<sup>13-15</sup>, whose  
86 working principle has been validated and reported previously<sup>16-18</sup>. We then performed  
87 experiments with HTTExon1-Q25 or buffer alone to exclude non-specific signals, and  
88 identified two compounds that interact with both LC3B and mHTTExon1-Q72, but not  
89 HTTExon1-Q25: 10O5 (GW5074,  
90 3-3-[(3,5-Dibromo-4-hydroxyphenyl)methylidene]-5-iodo-1H-indol-2-one) and 8F20  
91 (ispinesib,  
92 N-(3-aminopropyl)-N-[(1R)-1-[7-chloro-4-oxo-3-(phenylmethyl)-2-quinazolinyl]-2-meth

93 ylpropyl]-4-methylbenzamide) (Fig. 1b, annotated based on their ID in the compound  
94 library). We then measured the  $K_{on}$  and  $K_{off}$  of these interactions to confirm our  
95 observation (Fig. 1c-d) and revealed ~100 nM  $K_d$  of the compounds' interaction with  
96 LC3B or mHTT<sub>exon1-Q72</sub>. As shown in Fig. 1e, these compounds also interacted  
97 with the full-length mHTT (flHTT-Q73, Extended Data Fig. 1e), but not wtHTT  
98 (HTT<sub>exon1-Q25</sub> or flHTT-Q23, Fig. 1c&e) or irrelevant proteins (Extended Data Fig.  
99 2a) including MBP-His8 (MBP), superfolder GFP (sfGFP), and Rpn10 (a proteasomal  
100 ubiquitin-receptor) (Extended Data Fig. 1f). We then validated the interaction by an  
101 orthogonal assay, microscale thermophoresis (MST), and obtained consistent results  
102 (Extended Data Fig. 2b).

103 *mHTT-LC3 linker compounds reduced mHTT levels via autophagy in an*  
104 *allele-selective manner*

105 We then tested if these potential mHTT-LC3 linker compounds decrease mHTT  
106 levels *via* autophagy as predicted. Both hits decreased levels of mHTT in cultured  
107 primary cortical neurons from a well-established HD knockin mouse model  
108 (Hdh<sup>Q7/Q140</sup>)<sup>19</sup> (Fig. 2a), while had little or no effect on levels of wtHTT in the  
109 heterozygous HD neurons (Hdh<sup>Q7/Q140</sup>) (Fig. 2a) and wild-type neurons (Hdh<sup>Q7/Q7</sup>) (Fig.  
110 2b), consistent with their lack of wtHTT interaction. We then looked for other  
111 mHTT-LC3 linker compounds based on the common features of the two hit  
112 compounds 10O5 and 8F20. The –OH group in 10O5 and the –NH<sub>2</sub> group in 8F20  
113 were utilized in the nucleophile-isocyanate reaction for stamping of the SMMs, and  
114 these groups were inaccessible to mHTT and LC3B for the compound-protein

115 interaction during the screening (Fig. 2c). Thus, while these two hit compounds have  
116 different structures, their exposed chemical groups on the SMMs shared similarities  
117 by containing an aryl ring connected with a lactam-based bicyclic structure with  
118 halogen-substituted aryl group (Fig. 2c). We tested several compounds with similar  
119 features and identified two additional mHTT-LC3 linker compounds (Fig. 2c, AN1  
120 (3-5-Bromo-3-[(3-bromo-4,5-dihydroxyphenyl)methylidene]-1H-indol-2-one) and AN2  
121 (5,7-Dihydroxy-4-phenylcoumarin), for analog1 and analog2, respectively) that  
122 interact with both mHTT and LC3B, but not wtHTT or irrelevant control proteins  
123 (Extended Data Fig. 2c-d). They also reduced the levels of mHTT in an  
124 allele-selective manner in cultured HD mouse neurons (Fig. 2d). No cytotoxicity was  
125 observed in cultured neurons treated with these compounds at the tested  
126 concentration range (Extended Data Fig. 2e), confirming that the mHTT lowering was  
127 not due to cell loss.

128 Most of these compounds showed an optimal dose (hook effect) in lowering  
129 mHTT (Fig. 2a&d): a sufficient concentration is desired for tethering mHTT and LC3  
130 together, but excessively high concentrations may cause the compound molecules to  
131 interact with mHTT and LC3 separately, without tethering them. Similar concentration  
132 dependent effects were observed in HD patient fibroblasts (Fig. 3c, *right* panel) and  
133 have been reported for PROTAC<sup>20</sup>. Consistent with the prediction that the mHTT  
134 lowering is mediated by degradation *via* autophagy, the autophagy inhibitor NH<sub>4</sub>Cl or  
135 chloroquine blocked the mHTT-lowering effects (Extended Data Fig. 3a), confirming  
136 that the compounds targeted mHTT for autophagic degradation. Meanwhile, the

137 compound-induced mHTT-lowering effects were only slightly enhanced by the mTOR  
138 inhibitor rapamycin, an enhancer of autophagosome formation (Extended Data Fig.  
139 3a, *right* panels; also see Fig. 3b).

140 The mHTT lowering could be detected by multiple mHTT antibodies, including  
141 3B5H10, which detects a toxic species of the expanded polyQ stretch<sup>21,22</sup> (Fig. 2d,  
142 *right* panel), suggesting that the detected mHTT lowering was not due to affinity  
143 changes to a specific antibody. In addition, we did not observe any obvious increase  
144 of possible polyQ-containing mHTT fragments at lower molecular weights (Extended  
145 Data Fig. 3b-c), suggesting that the mHTT lowering was not due to increased  
146 site-specific cleavages of mHTT.

147 We further investigated the compounds' effects in HD patient cells by the  
148 well-established HTRF (Homologous Time-Resolved Fluorescence) assay<sup>23,24</sup>, which  
149 is more quantitative than Western-blot, although inapplicable to mouse mHTT  
150 proteins due to non-specific signals<sup>25</sup>. We observed autophagy-dependent lowering of  
151 mHTT by these compounds in HD patient fibroblasts and iPSC-derived neurons (Fig.  
152 3a-b&Extended Data Fig. 3d), but not wtHTT in the wild-type or Parkinson's disease  
153 (PD) patient fibroblasts (Fig. 3a). To further confirm the role of autophagic degradation,  
154 we tested the compounds' effect with or without lowering of ATG5, a key autophagy  
155 gene required for autophagosome formation<sup>26</sup>. ATG5 knockdown in HD patient  
156 fibroblasts (Q47) significantly decreased LC3-II and nullified the mHTT lowering  
157 effects induced by the mHTT-LC3 linker compounds (Extended Data Fig. 3e). Similar  
158 results were obtained in ATG5 knockout mouse embryonic fibroblasts<sup>26</sup> (MEFs,

159 Extended Data Fig. 3f), confirming that the compounds' effects were mediated by  
160 autophagic degradation.

161 The two hit compounds 10O5 and 8F20 are known to inhibit c-Raf and KSP<sup>27,28</sup>,  
162 respectively, whereas AN1 and AN2 had unknown activities on these targets. We thus  
163 tested their potential influences on c-Raf and KSP. Based on the *in vitro* c-Raf kinase  
164 assay, 10O5, the known c-Raf inhibitor, but not the other three compounds inhibited  
165 c-Raf at the concentrations tested (Extended Data Fig. 4a). We then tested Mek/Erk  
166 phosphorylation levels in the cultured neurons treated with these compounds at  
167 optimal mHTT-lowering concentrations to evaluate Raf activity<sup>29</sup> and found no  
168 significant effects of all tested compounds (Extended Data Fig. 4b, *left* panel). We  
169 also tested phospho-BUBR1 levels to evaluate KSP activities<sup>30</sup>, and observed no  
170 significant effects either (Extended Data Fig. 4b, *right* panel). We made similar  
171 observations in the HD patient fibroblasts (Q47) (Extended Data Fig. 4c). Thus, the  
172 observed mHTT-lowering is probably irrelevant to c-Raf or KSP inhibition. To further  
173 confirm this, we examined the effects of several known c-Raf or KSP inhibitors, and  
174 found that they had no HTT-lowering effects (Fig. 3c, *left* panel). Two of these  
175 inhibitors, PLX-4720 and BAY1217389, have structures somewhat similar to 10O5  
176 and 8F20, respectively (Fig. 3c, *middle* panel). They showed no effects in lowering  
177 mHTT in patient cells at sub-micromolar concentrations (Fig. 3c, *right* panel), probably  
178 because they had very weak affinity to LC3 and mHTT, if any (Extended Data Fig. 2c,  
179 *right panels*). In comparison, AN2 showed a dose-dependent mHTT lowering in the  
180 same cells (Fig. 3c, *right* panel).

181 We then investigated the compounds' effects *in vivo*. Since the *Drosophila* LC3  
182 homolog Atg8 has a predicted structure highly similar to LC3B (Extended Data Fig.  
183 5a), we tested the compounds in HD transgenic flies expressing human full-length  
184 mHTT. All of the discovered mHTT-LC3 linker compounds significantly reduced mHTT  
185 levels (Extended Data Fig. 5b), validating the *in vivo* efficacy of these compounds.

186 We further investigated the compounds' effects *in vivo* in the HD knockin mouse  
187 model (Hdh<sup>Q7/Q140</sup>)<sup>19</sup> by intracerebroventricular (icv) injections. Three out of the four  
188 linker compounds (10O5, AN1 or AN2, but not 8F20) led to significant lowering of  
189 mHTT in cortices of HD mice (Extended Data Fig. 6a). We then performed  
190 intraperitoneal (ip) injection of 10O5 and AN2 at 0.5 mg/kg in HD knockin mice. The  
191 compounds crossed the blood-brain barrier and reached the brain at detectable  
192 concentrations (Extended Data Fig. 5c, ~20 to 200 nM for 10O5 and ~20 to 40 nM for  
193 AN2; no signal was detected in the DMSO injected control group) 0.5 ~ 6 hours after  
194 injection. Consistent with this, we observed significant allele-selective lowering of  
195 mHTT in mouse cortices and striata (Extended Data Fig. 6b-c). The observed  
196 lowering was not due to changes in mHTT solubility, because no increase of mHTT  
197 aggregates was observed in the cortical tissues of mice treated with these  
198 compounds (Extended Data Fig. 6d).

#### 199 *mHTT-LC3 linker compounds tethered mHTT to autophagosomes*

200 We then examined whether these compounds truly function as linkers between  
201 mHTT and LC3 to target mHTT for autophagosome engulfment. Presence of 10O5 or  
202 AN2, the two compounds that were effective *in vivo* by ip-injection, markedly

enhanced the mHTT-LC3 interaction in the *in vitro* pull-down experiments (Fig. 4a, comparing lane 11 with 12&13; Fig. 4b, comparing lane 11 with 12 for both *upper* and *lower* panels). No enhancement effect was observed for the wtHTT-LC3 interaction (Fig. 4a, lane 8-10; Fig. 4b, lane 9-10 in both *upper* and *lower* panels). Consistent with this, these compounds led to elevated engulfment of mHTT by autophagosomes both in transiently transfected HeLa cells expressing exogenous GFP-LC3B and HTTexon1-MBP-His fragments (Fig. 4c), and in mouse striatal cells (STHdh<sup>Q111/Q111</sup>)<sup>31</sup> expressing endogenous LC3 and full-length mHTT proteins (Fig. 4d).

The data above confirmed that the compounds tethered mHTT and LC3B/autophagosomes *in vitro* and in cells, although the detailed structural information remains to be resolved.

*mHTT-LC3 linker compounds did not influence autophagy function and reduced mHTT in a relatively specific manner*

The mHTT-lowering by the linker compounds was unlikely due to an enhancement of autophagy functions, because the number and size of autophagosomes remained unchanged (Extended Data Fig. 7a). We then further investigated whether the compounds could influence autophagy by established approaches in the literature<sup>32-34</sup>. Neither 10O5 nor AN2 influenced the autophagosome-lysosome fusion or autophagy activity (Extended Data Fig. 7b-d). Furthermore, we observed no changes of the LC3-II level in the cultured cortical neurons treated with 10O5 or AN2 in the absence or presence of the lysosome inhibitor bafilomycin-A1 (bafA1) (Extended Data Fig. 7e). The level of the known

225 autophagy-selective substrate protein SQSTM1/p62 was also unaffected *in vivo* and  
226 in cultured neurons (Extended Data Fig. 7f & 8a). In addition, other wild-type polyQ  
227 proteins (Atxn3 and Tbp) and control proteins (Nbr1, Ncoa4, Actin, Gapdh, Tubulin)  
228 were not influenced (Extended Data Fig. 8a, all changes < 10%).

229 We then performed proteomics analysis to obtain a more complete scope of  
230 proteins that may have been influenced. We observed significant lowering (~20%,  $p <$   
231 0.01) of HTT levels in cortices of mice ip-injected with 10O5 or AN2 (Extended Data  
232 Fig. 8b, bar plots). Considering that the proteomics analysis was unable to distinguish  
233 mHTT from wtHTT, the actual mHTT lowering was expected to be more. Meanwhile,  
234 using the criteria of  $p < 0.01$ , we observed changes of only a small percentage of  
235 proteins (Extended Data Fig. 8b, and see Supplementary Table 2 for details). No  
236 autophagy-specific substrate proteins exhibited significant changes and there was no  
237 enrichment of autophagy pathway-related proteins (Supplementary Table 2), further  
238 confirming that autophagy was unaffected. Proteomics analysis in cultured neurons  
239 revealed consistent results (Extended Data Fig. 8c, and see Supplementary Table 3  
240 for details).

#### 241 *mHTT-LC3 linker compounds lowered other proteins with expanded polyQ*

242 The discovered linker compounds interacted with and lowered mHTT but not  
243 wtHTT (Fig. 1). The simplest explanation is that the compounds specifically interact  
244 with the expanded polyQ tract, possibly by recognizing its emergent conformation that  
245 is different from the short polyQ stretch<sup>21,35</sup>. If so, the discovered linker compounds  
246 may reduce other proteins with expanded polyQ. Consistent with this prediction, the



linker compounds 10O5, AN1 and AN2 reduced the levels of mutant but not wild-type ATXN3 in SCA3 patient fibroblasts (Extended Data Fig. 9a) and exogenously expressed 72Q-GFP, 46Q-GFP, 38Q-GFP but not 25Q-GFP proteins (containing just the Met-polyQ-sfGFP sequences) in HEK293T cells (Extended Data Fig. 9b). These data suggest that the compounds distinguished the expanded from the short polyQ stretch at a threshold between 25Q and 38Q. To further confirm this, we tested the compound-polyQ interactions (Extended Data Fig. 9c), and confirmed that 10O5, AN1 and AN2 interacted with polyQ-GFP with 38Q or longer polyQ, but not 25Q-GFP or GFP alone (Extended Data Fig. 9d-e; Extended Data Fig. 2).

#### *mHTT-LC3 linker compounds rescued HD-relevant phenotypes*

We further investigated the therapeutic potential of the compounds in treating HD. All the mHTT-LC3 linker compounds rescued mHTT toxicity in HD patient iPSC-derived neurons (Fig. 5a-b). They also rescued HD-relevant behavioral deficits and increased the lifespan of flies expressing human mHTT, while having no influence on the flies expressing wtHTT (Fig. 5c).

Finally, we investigated the disease-relevant behavioral phenotypes in 10-months old heterozygous HD knockin mouse ( $Hdh^{Q7/Q140}$ ). Significant deficits of HD mice were observed in several behavioral tests including rotarod, balance beam, and gripping force tests, while the DMSO (110  $\mu$ g/kg) injection alone did not have an effect (Extended Data Fig. 9f-h). In comparison, ip injection of the 10O5 or AN2 significantly improved HD-relevant behavioral deficits in these tests, without influencing the wild-type mice (Fig. 5d-f), demonstrating a rescue of HD-relevant phenotypes. This is

269 a proof of principle study, and further investigations will be required to establish the  
270 suitability for therapeutic application.

## 271 **Discussion**

272 In this study, we have identified mHTT-LC3 linker compounds that were able to  
273 reduce mHTT levels at ~nM concentrations in HD cells and at 0.5 mg/kg by ip  
274 injection *in vivo* (Extended Data Table 1). The compounds did not influence wtHTT,  
275 which has essential functions especially during development and young adulthood<sup>36</sup>.  
276 These features of the compounds are highly desired for HD treatment or potentially  
277 other polyQ diseases (Extended Data Fig. 9a-e), although preclinical studies of  
278 longitudinal efficacy and safety are necessary for therapeutic development.

279 From a broader perspective, we demonstrate the concept of using small molecule  
280 compounds to “glue” the target protein (mHTT) and LC3 for autophagic degradation  
281 (Fig. 5g). We selected mHTT as our target protein, because wtHTT provides a perfect  
282 control for screening. We had the fortune to discover hits that interact with both LC3B  
283 and mHTT. If no such hits had been identified, linker compounds could still be  
284 generated by conjugating an mHTT-interacting compound and an LC3-interacting  
285 compound by the nucleophile-isocyanate reaction utilized for the stamping of SMMs.  
286 To develop this concept, the critical next step is to resolve the core chemical  
287 compartment that interacts with LC3 without influencing its function. Comprehensive  
288 medicinal chemistry and structural studies are needed to resolve the compound-LC3  
289 interaction interface, which may then be developed to a general degradation tool for  
290 conjugation with other compounds interacting with specific targets of interest.

291 In summary, we have identified mHTT-LC3 linker compounds that are capable of  
 292 lowering mHTT *in vivo* in an allele-selective manner, and demonstrated the possibility  
 293 of utilizing autophagosome-tethering compounds (ATTEC) to lower target proteins,  
 294 providing new entry points for drug discovery.

295

## 296 Reference

- 297 1 Kabeya, Y. *et al.* LC3, A mammalian homologue of yeast Apg8p, is localized in  
 298 autophagosome membranes after processing. *Embo J* **19**, 5720-5728, doi:  
 299 10.1093/emboj/19.21.5720 (2000).
- 300 2 Scherzinger, E. *et al.* Huntingtin-encoded polyglutamine expansions form  
 301 amyloid-like protein aggregates in vitro and in vivo. *Cell* **90**, 549-558, doi:  
 302 10.1016/s0092-8674(00)80514-0 (1997).
- 303 3 Warrick, J. M. *et al.* Expanded polyglutamine protein forms nuclear inclusions  
 304 and causes neural degeneration in Drosophila. *Cell* **93**, 939-949,  
 305 doi:10.1016/s0092-8674(00)81200-3 (1998).
- 306 4 Fire, A. *et al.* Potent and specific genetic interference by double-stranded RNA  
 307 in *Caenorhabditis elegans*. *Nature* **391**, 806-811, doi:10.1038/35888 (1998).
- 308 5 Mali, P. *et al.* RNA-guided human genome engineering via Cas9. *Science* **339**,  
 309 823-826, doi:10.1126/science.1232033 (2013).
- 310 6 Cong, L. *et al.* Multiplex genome engineering using CRISPR/Cas systems.  
 311 *Science* **339**, 819-823, doi:10.1126/science.1231143 (2013).
- 312 7 Winter, G. E. *et al.* Phthalimide conjugation as a strategy for in vivo target  
 313 protein degradation. *Science* **348**, 1376-1381, doi:10.1126/science.aab1433  
 314 (2015).
- 315 8 Lu, K., den Brave, F. & Jentsch, S. Pathway choice between proteasomal and  
 316 autophagic degradation. *Autophagy* **13**, 1799-1800,  
 317 doi:10.1080/15548627.2017.1358851 (2017).
- 318 9 Mizushima, N., Levine, B., Cuervo, A. M. & Klionsky, D. J. Autophagy fights  
 319 disease through cellular self-digestion. *Nature* **451**, 1069-1075,  
 320 doi:10.1038/nature06639 (2008).
- 321 10 Zhu, C. *et al.* Developing an efficient and general strategy for immobilization of  
 322 small molecules onto microarrays using isocyanate chemistry. *Sensors (Basel)*  
 323 **16**, doi:10.3390/s16030378 (2016).
- 324 11 Fei, Y. *et al.* Screening small-molecule compound microarrays for protein  
 325 ligands without fluorescence labeling with a high-throughput scanning  
 326 microscope. *J Biomed Opt* **15**, 016018, doi:10.1117/1.3309743 (2010).
- 327 12 Mangiarini, L. *et al.* Exon 1 of the HD gene with an expanded CAG repeat is  
 328 sufficient to cause a progressive neurological phenotype in transgenic mice.  
 329 *Cell* **87**, 493-506, doi:S0092-8674(00)81369-0 (1996).

330 13 Liu, H. *et al.* Nuclear cGAS suppresses DNA repair and promotes  
331 tumorigenesis. *Nature* **563**, 131-136, doi:10.1038/s41586-018-0629-6 (2018).

332 14 Landry, J. P. *et al.* Discovering small molecule ligands of vascular endothelial  
333 growth factor that block VEGF-KDR binding using label-free microarray-based  
334 assays. *Assay Drug Dev Technol* **11**, 326-332, doi:10.1089/adt.2012.485  
335 (2013).

336 15 Fei, Y. *et al.* Characterization of receptor binding profiles of influenza A viruses  
337 using an ellipsometry-based label-free glycan microarray assay platform.  
338 *Biomolecules* **5**, 1480-1498, doi:10.3390/biom5031480 (2015).

339 16 Zhu, X. *et al.* Oblique-incidence reflectivity difference microscope for label-free  
340 high-throughput detection of biochemical reactions in a microarray format.  
341 *Appl Opt* **46**, 1890-1895, doi:10.1364/ao.46.001890 (2007).

342 17 Landry, J. P., Zhu, X. D. & Gregg, J. P. Label-free detection of microarrays of  
343 biomolecules by oblique-incidence reflectivity difference microscopy. *Opt Lett*  
344 **29**, 581-583, doi:10.1364/ol.29.000581 (2004).

345 18 Zhu, C. *et al.* Fast focal point correction in prism-coupled total internal  
346 reflection scanning imager using an electronically tunable lens. *Sensors*  
347 (*Basel*) **18**, doi:10.3390/s18020524 (2018).

348 19 Menalled, L. B., Sison, J. D., Dragatsis, I., Zeitlin, S. & Chesselet, M. F. Time  
349 course of early motor and neuropathological anomalies in a knock-in mouse  
350 model of Huntington's disease with 140 CAG repeats. *J Comp Neurol* **465**,  
351 11-26, doi:10.1002/cne.10776 (2003).

352 20 Bondeson, D. P. *et al.* Catalytic in vivo protein knockdown by small-molecule  
353 PROTACs. *Nat Chem Biol* **11**, 611-617, doi:10.1038/nchembio.1858 (2015).

354 21 Miller, J. *et al.* Identifying polyglutamine protein species in situ that best predict  
355 neurodegeneration. *Nat Chem Biol* **7**, 925-934, doi:10.1038/nchembio.694  
356 (2011).

357 22 Fu, Y. *et al.* A toxic mutant huntingtin species is resistant to selective  
358 autophagy. *Nat Chem Biol* **13**, 1152-1154, doi:10.1038/nchembio.2461 (2017).

359 23 Baldo, B. *et al.* TR-FRET-based duplex immunoassay reveals an inverse  
360 correlation of soluble and aggregated mutant huntingtin in huntington's  
361 disease. *Chem Biol* **19**, 264-275, doi:10.1016/j.chembiol.2011.12.020 (2012).

362 24 Weiss, A. *et al.* Single-step detection of mutant huntingtin in animal and  
363 human tissues: a bioassay for Huntington's disease. *Anal Biochem* **395**, 8-15,  
364 doi:10.1016/j.ab.2009.08.001 (2009).

365 25 Lu, B. *et al.* Identification of NUB1 as a suppressor of mutant Huntington  
366 toxicity via enhanced protein clearance. *Nat Neurosci* **16**, 562-570,  
367 doi:10.1038/nn.3367 (2013).

368 26 Mizushima, N. *et al.* Dissection of autophagosome formation using  
369 Apg5-deficient mouse embryonic stem cells. *J Cell Biol* **152**, 657-668,  
370 doi:10.1083/jcb.152.4.657 (2001).

371 27 Lackey, K. *et al.* The discovery of potent cRaf1 kinase inhibitors. *Bioorg Med*  
372 *Chem Lett* **10**, 223-226, doi:10.1016/s0960-894x(99)00668-x (2000).

373 28 Reddy, K. & D'Orazio, A. Highlights from the international conference on

374 molecular targets and cancer therapeutics: discovery, biology, and clinical  
375 applications, Philadelphia, PA. ECCO 13--The European Cancer Conference,  
376 Paris, France, October 30-November 3, 2005. *Clin Genitourin Cancer* **4**,  
377 156-159, doi:10.1016/S1558-7673(11)70136-7 (2005).

378 29 Johnson, G. L. & Lapadat, R. Mitogen-activated protein kinase pathways  
379 mediated by ERK, JNK, and p38 protein kinases. *Science* **298**, 1911-1912,  
380 doi:10.1126/science.1072682 (2002).

381 30 Tao, W. *et al.* An inhibitor of the kinesin spindle protein activates the intrinsic  
382 apoptotic pathway independently of p53 and de novo protein synthesis. *Mol*  
383 *Cell Biol* **27**, 689-698, doi:10.1128/MCB.01505-06 (2007).

384 31 Trettel, F. *et al.* Dominant phenotypes produced by the HD mutation in  
385 STHdh(Q111) striatal cells. *Hum Mol Genet* **9**, 2799-2809,  
386 doi:10.1093/hmg/9.19.2799 (2000).

387 32 Kimura, S., Noda, T. & Yoshimori, T. Dissection of the autophagosome  
388 maturation process by a novel reporter protein, tandem fluorescent-tagged  
389 LC3. *Autophagy* **3**, 452-460, doi:10.4161/auto.4451 (2007).

390 33 Zhang, J., Wang, J., Ng, S., Lin, Q. & Shen, H. M. Development of a novel  
391 method for quantification of autophagic protein degradation by AHA labeling.  
392 *Autophagy* **10**, 901-912, doi:10.4161/auto.28267 (2014).

393 34 Ni, H. M. *et al.* Dissecting the dynamic turnover of GFP-LC3 in the  
394 autolysosome. *Autophagy* **7**, 188-204, doi:10.4161/auto.7.2.14181 (2011).

395 35 Feng, X., Luo, S. & Lu, B. Conformation polymorphism of polyglutamine  
396 Proteins. *Trends Biochem Sci* **43**, 424-435, doi:10.1016/j.tibs.2018.03.002  
397 (2018).

398 36 Wang, G., Liu, X., Gaertig, M. A., Li, S. & Li, X. J. Ablation of huntingtin in adult  
399 neurons is nondeleterious but its depletion in young mice causes acute  
400 pancreatitis. *Proc Natl Acad Sci U S A* **113**, 3359-3364,  
401 doi:10.1073/pnas.1524575113 (2016).

402 37 Vijayvargia, R. *et al.* Huntingtin's spherical solenoid structure enables  
403 polyglutamine tract-dependent modulation of its structure and function. *Elife* **5**,  
404 e11184, doi:10.7554/eLife.11184 (2016).

405

## Main figure legends

### Figure 1. Identification of potential mHTT-LC3 linker compounds by SMM-based screening and validation

a) The OI-RD image of a small molecule microarray (SMM). Each compound was printed in duplicates in adjacent vertical positions.

b) Magnified view of surface mass density changes after incubation with HTTexon1-Q25-MBP, HTTexon1-Q72-MBP, and LC3B, respectively. Two hits (10O5 and 8F20) were highlighted by red rectangular boxes.

c-e) Association-dissociation curves of surface immobilized compounds 8F20 and 10O5 with HTTexon1-Q72-MBP (Q72), HTTexon1-Q25-MBP (Q25), LC3B, full-length HTT-Q73 (Q73), or full-length HTT-Q23 (Q23) at the indicated concentrations of purified proteins. For details see "Methods".

### Figure 2. mHTT-LC3 linker compounds lowered mHTT but not wtHTT through autophagy in cultured mouse neurons.

a) Western-blots (HTT detected by the 2166 antibody) and quantifications of compound-treated cultured cortical neurons from Hdh<sup>Q7/Q140</sup> HD knockin mice. The statistical analysis was performed by two-way ANOVA tests. For 10O5,  $F(1, 72) = 50.93$ ,  $p < 0.0001$ ; for 8F20,  $F(1, 40) = 8.903$ ,  $p = 0.0048$ .

b) Representative Western-blots (from 3 biological repeats) of cultured wild-type cortical neurons treated with the indicated compounds.

c) 2D-structures of the hit compounds and the other effective linker compounds identified. The red lines indicate the glass-chips used to stamp the hit compounds. The dotted ovals indicate the possible chemical groups exposed for protein-compound interactions in the screening.

d) *Left* and *middle* panels: similar to a), but treated with the compounds AN1 or AN2. For AN1,  $F(1, 70) = 32.96$ ,  $p < 0.0001$ ; For AN2,  $F(1, 69) = 23.03$ ,  $p < 0.0001$ . *Right* panels: similar to a), but blotted with different HTT antibodies (10O5 and 8F20: 100nM; AN2: 50 nM).

For all panels, n indicates the number of independently plated wells, and bars represent mean and s.e.m.. Full-blots of cropped gels are shown in Extended Data Fig. 3b or Supplementary Fig. 1.

### Figure 3. mHTT-LC3 linker compounds lowered mHTT in HD patient cells.

a) HTT levels measured by HTRF (2B7/MW1 for mHTT, and 2B7/2166 for total HTT) in the indicated HD, WT or PD patient primary fibroblasts treated with the indicated compounds (100 nM). All signals were normalized to the averaged signals from the DMSO control group. The statistical analysis was performed by one-way ANOVA with post-hoc Dunnett's tests. "\*\*\*\*":  $p < 0.0001$ .

b) Similar to a), but in immortalized HD patient fibroblasts treated with or without the autophagy inhibitors NH<sub>4</sub>Cl/chloroquine/bafilomycin-A1, or the autophagy enhancer rapamycin.

c) *Left* panel: similar to a), but in immortalized HD patient fibroblasts (Q47) treated with indicated c-Raf or KSP inhibitors at 100 nM. *Middle* panel: 2D structure of

inhibitors. The dotted ovals indicate the parts of the compounds that share similarities with the hit compounds. *Right* panel: dose-dependent curves of the indicated compounds.

For all panels, n indicates the number of independently plated wells, and bars represent mean and s.e.m..

**Figure 4. Linker compounds enhanced mHTT-LC3 interaction and tethered mHTT to autophagosomes**

a-b) Representative results (from 3 biological repeats) of *in vitro* pull-down experiments using purified HTT and LC3B proteins. For details see “Methods”.

c-d) Representative images (*scale* bar: 10  $\mu$ m) and quantifications of the co-localization between HTT and autophagosomes. For details see “Methods”.

Bars present mean and s.e.m.. The n number indicates the number of cells. The statistical analysis was performed by one-way ANOVA with post-hoc Dunnett’s tests.

“\*\*\*\*”:  $p < 0.0001$ .

**Figure 5. Linker compounds rescued HD-relevant phenotypes in cells and *in vivo*.**

a) Representative immunostaining images (*scale* bar: 50  $\mu$ m) and quantifications of the neuronal specific tubulin marker TUBB3 and DAPI staining showing neuronal morphology of patient iPSC-derived striatal neurons (HD: Q47; WT: Q19) treated with indicated compounds. For details see “Methods”.

b) Neuronal apoptosis measurement at different time points after BDNF removal using a green fluorescent dye (NucView 488) detecting active caspase-3. For details see “Methods”.

c) *Left*: Kaplan-Meier survival curves of transgenic *Drosophila* with the indicated transgenes and compound treatments. *Right*: similar to the *left* panel, but plotting the climbing performance as a function of age after eclosion. For details see “Methods”.

d-f) Mouse behavioral tests showing the improvement of HD-relevant phenotypes by ip-injection of the indicated compounds at 0.5 mg/kg. For details see “Methods”.

g) A schematic model showing how mHTT-LC3 linker compounds may induce mHTT degradation, illustrating the concept of lowering target proteins by autophagosome-tethering compounds (ATTEC). The images representing HTT proteins were published previously<sup>37</sup>.

## Methods

### Additional details of figure legends

Figure 1. Identification of potential mHTT-LC3 linker compounds by SMM-based screening and validation.

c-e) For all association-dissociation curves, vertical dash lines mark the starts of association and dissociation phases of the binding event. The red dash curves are global fits to a Langmuir reaction model with the fitting parameters listed at the bottom of each plot. No binding signals were observed for HTTexon1-Q25-MBP or full-length HTT-Q23 proteins, and thus the parameters were not presented.

Figure 4. Linker compounds enhanced mHTT-LC3 interaction and tethered mHTT to autophagosomes.

a-b) For MBP pull-down (a), purified HTTexon1-MBP (10 µg) or MBP (10 µg) bound MBP-resin were incubated with the purified LC3B protein (40 µg) and the indicated compounds. The HTTexon1-MBP or the MBP proteins were pulled down and the eluates were tested for co-precipitated LC3B. 4% of the total eluate was loaded in each lane, and the input : pull-down loading ratio was 100%. Both 10O5 and AN2 enhanced LC3B's interaction with HTTexon1-Q72-MBP, but not HTTexon1-Q25-MBP. Note that the MBP blot signals were much weaker for the Q72 protein, possibly because recognition of the MBP tag by the antibody was affected in the fusion protein. Meanwhile, data interpretation was not influenced, because compound treatments did not alter the MBP signals for the Q72 protein (last three lanes). The GST pull-down (b) was performed similarly, except using full-length HTT-Q73 or full-length HTT-Q23 (both without fusion tags) and GST-LC3B proteins for the *in vitro* GST pull-down experiments to precipitate GST-LC3B or GST alone with its binding proteins, and then eluted for detection. Note that the pull-down is in the reverse direction of the pull-down in (a). The input : pull-down loading ratio for the GST blot was 100%, whereas the ratio for the HTT blot was 10% to avoid over-exposure of the input. Both 10O5 and AN2 enhanced LC3B's interaction with the full-length HTT-Q73 but not the full-length HTT-Q23 protein.

c-d) Representative confocal microscopy images (*scale bar*: 10 µm) and quantifications of the co-localization between HTTexon1-MBP-His (red, detected by anti-His immunofluorescence) and LC3B-GFP (green, detected by GFP fluorescence directly) in transiently transfected HeLa cells (c) or between endogenous mHTT and LC3-II in the HD knockin mouse striatal cells (STHdh<sup>Q111/Q111</sup>) (d). For over-expressed proteins (c), the LC3B-GFP alone transfected or the HTTexon1-MBP-His alone transfected cells were imaged at both channels to ensure the specificity of the signals (*upper panels*). The white arrows indicate representative co-localization puncta. Parts



of the images have been amplified to show co-localization puncta more clearly (pointed by orange arrows). Since the puncta were obvious, co-localization was analyzed by counting the red<sup>+</sup>green<sup>+</sup> (yellow) and the total red<sup>+</sup> puncta directly, and then calculate their ratio for each cell. Blind analysis was performed for quantifications. For endogenous proteins (d), mHTT was detected by the anti-HTT antibody 2166, and the endogenous LC3-II was detected by an anti-LC3 antibody that has been reported to specifically detect LC3-II<sup>32</sup>. Since the signals of endogenous proteins were more dispersed, the co-localization analysis was performed blindly by measuring the red<sup>+</sup>green<sup>+</sup> (yellow) and the total red<sup>+</sup> pixels by ImageJ, and then calculate their ratio for each cell.

Figure 5. Linker compounds rescued HD-relevant phenotypes in cells and *in vivo*.

a) Loss of processes and shrinkage of neurons were observed in HD neurons after BDNF removal. Bar plots: quantification of the TUBB3 signal covered area (TUBB3 area) normalized to the nuclei counts based on DAPI. The lower TUBB3 area *per cell* reflects neuronal processes shrinkage and loss. Data were normalized to the average of WT controls. The statistical analysis was performed by one-way ANOVA (F (5, 60) = 94.78) with post-hoc Dunnett's tests: "\*\*\*\*": p<0.0001. The n number indicates the number of independently plated wells.

b) The images were captured every 3 hours inside the incubator using Incucyte, and the caspase-3 active cells were quantified by the fluorescent object count per field. The statistical analysis was performed by two-way ANOVA (F (43, 516) = 12.85) with post-hoc Dunnett's tests, comparing to the HD\_DMSO group. "\*\*\*\*": p<0.0001. The numbers in brackets indicate the number of independently plated wells, with 4 fields *per well* imaged and averaged for quantification. Three batches were tested and showed consistent results.

c) *Left*: the *Drosophila* expressed full-length HTT proteins (Q128 or Q16) in the nervous system driven by *elav-GAL4*. 75 flies were tested for each group. The statistics was performed by Log-rank (Mantel-Cox) test, comparing compound treated groups with DMSO controls in Q128 flies. \*\*\*\*: p < 0.0001. *Right*: similar as the *left* panel, but plotting the climbing performance as a function of age after eclosion. The statistics was performed by two-way ANOVA (F (4, 275) = 122.1) with post-hoc Dunnett's tests, comparing the compound treated groups with the DMSO controls in Q128 flies. Numbers in brackets indicate the number of vials (each containing 15 flies) tested. \*\*\*\*: p < 0.0001.

d-f) The numbers in brackets indicated the number of mice tested. The statistical analysis was performed by two-way ANOVA with post-hoc Dunnett's tests, and the p

559 values compared with the DMSO control were indicated (\*\*\*\*:  $p < 0.0001$ ). For HD  
560 mice,  $F(2, 195) = 4.963$  in rotarod tests,  $F(2, 195) = 37.31$  in balance beams tests,  
561 and  $F(2, 156) = 7.068$  in gripping force tests. No significant difference was detected in  
562 the wild-type mice injected with different compounds. The investigators were blinded  
563 to the compounds and genotypes when performing the experiments. In all panels, the  
564 error bars represent mean and s.e.m..

#### 565 **Compound stamping on the microarray**

566 Small molecule microarrays (SMMs) containing 3,375 bioactive compounds were  
567 used for high-throughput screening of target proteins. The compound library  
568 containing 1,527 drugs approved by Food and Drug Administration (FDA) of United  
569 States, 1,053 natural products from traditional Chinese medicine, and 795 known  
570 inhibitors were stamped onto the SMMs. Each compound was dissolved in DMSO at  
571 a concentration of 10 mM and printed in duplicates along vertical direction on  
572 homemade phenyl-isocyanate functionalized glass slides with a contact microarray  
573 printer (SmartArrayer 136, CapitalBio Corporation). Biotin-BSA at a concentration of  
574 7,600 nM in 1× phosphate-buffered saline (PBS) and biotin-(PEG)<sub>2</sub>-NH<sub>2</sub> at a  
575 concentration of 5 mM in DMSO were printed as the inner and outer borders of SMMs,  
576 respectively. The diameter of each spot was about 150 μm and spacing between two  
577 adjacent spots was 250 μm. The printed SMMs were then dried at 45 °C for 24 h to  
578 facilitate covalent bonding of nucleophilic groups of small molecules to isocyanate  
579 groups of the functionalized slides. Afterwards, the SMMs were stored in a -20 °C  
580 freezer.

#### 581 **Expression and purification of recombinant proteins**

582 The human microtubule associated protein 1 light chain 3 beta (MAP1LC3B,  
583 LC3B) gene (GenBank: NM\_022818.4) was amplified by PCR and cloned into a  
584 pGEX-6P1 (GE Healthcare) derived vector pGHT, which is a prokaryotic expression  
585 vector reconstructed by adding a His8 tag and a TEV protease cleavage site before  
586 the pGEX-6P1 multiple cloning site. After sequencing verification, the expression  
587 plasmid pGHT-LC3B was introduced into *Escherichia coli* BL21 (DE3) pLysS, in which  
588 the recombinant GST-LC3B protein was expressed by IPTG induction. When the  
589 bacterial culture reached OD<sub>600</sub>=0.8, its temperature was decreased to 18 °C, and 0.2  
590 mM IPTG was added into the culture for an additional 20 h incubation. The cells were  
591 then harvested by centrifugation (6,000g, 4 °C, 15 min) and the cell pellet was  
592 suspended in 50 mM Tris-HCl buffer, pH 7.5, with 150 mM NaCl and 5% glycerol.  
593 Cells were then disrupted by sonication, followed by centrifugation (20,000g, 4 °C, 60  
594 min). The supernatants were then loaded onto a HisTrap HP column (GE Healthcare,

cat. no. 17524701), and eluted with 50 mM Tris-HCl buffer, pH 7.5, containing 150 mM NaCl, 5% glycerol and 300 mM imidazole. The LC3B eluate was then mixed with TEV protease (Sigma, cat. no. T4455; eluted protein: TEV protease = 100:1) and dialyzed against the dialysate buffer (50 mM Tris-HCl buffer, pH 7.5, containing 100 mM NaCl) in 4 °C overnight. After TEV protease treatment, the samples were then loaded onto a HisTrap HP column again, the flow through fraction which mainly contains tag removed recombinant LC3B. Afterwards, the proteins were concentrated and further purified by Superose 6 Increase 10/300 GL (GE Healthcare) size exclusive chromatography. Finally, the purified proteins were concentrated to approximately 10 mg/ml in 50 mM HEPES buffer with 100 mM NaCl for further analysis. The MBP-His8 and Rpn10 proteins were purified similarly.

The full-length HTT proteins, HTTexon1-MBP, polyQ-sfGFP and sfGFP proteins were purified from mammalian cells. For full-length HTT proteins, the human *HTT* gene (GenBank: NM\_002111.8) with (CAG)<sub>23</sub> or (CAG)<sub>73</sub> (23Q or 73Q for proteins) were *de novo* synthesized (by Genewiz Inc.), sequence validated and then cloned into a modified pCAG vector with an N-terminal protein A tag. The plasmid was transfected to human embryonic kidney E293 cells using polyethylenimine (PEI, from Polysciences, cat. no. 23966). After culture at 37 °C for 48 to 60 h, cells were collected and lysed at 4 °C for 1 h in lysis buffer containing 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 5% glycerol, 0.5% CHAPS, 3 mM DTT, 1% PMSF, 1 µg/ml pepstatin, 1 µg/ml leupeptin and 1 µg/ml aprotinin, 5 mM ATP and 5 mM MgCl<sub>2</sub>. After centrifugation at 15000 rpm for 40 min, the supernatants were then incubated with IgG monoclonal antibody-agarose (Smart-lifesciences, cat. no. SA030010) for 2 h and unbound proteins were extensively washed away. The HTT proteins were then digested using TEV protease overnight to remove the protein A tag and eluted protein was further purified by ion exchange and gel filtration chromatography using Mono Q and Superose 6 (5/150 GL) columns from GE healthcare. The peak fractions were pooled for further biochemical analysis. The HTTexon1 with 25Q or 72Q cDNA were also *de novo* synthesized and cloned into a mammalian expression vector pTT5SH8Q2 for large scale production in HEK293T cells. In order to improve the production yield and increase the solubility, a C terminal MBP tag was added after the HTTexon1 sequences to generate the pTT-HTTexon125Q-MBP and pTT-HTTexon125Q-MBP plasmids. For protein production and purification, the HEK293T cells were transfected by pTT-HTT25QExon1-MBP and pTT-HTT72QExon1-MBP plasmids with linear PEI (PolySciences #24765), and then collected after 48 h. The cells were then lysed by sonication in buffer containing 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 20 mM imidazole, 5% glycerol, protease inhibitor

cocktail (Sigma) and 50U/ml benzonase (Sigma). After centrifugation, the supernatants were loaded onto HisTrap HP column (GE Healthcare), and eluted with the buffer containing 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 300 mM imidazole, 5% glycerol and protease inhibitor cocktail. The MBP tag was not cleaved to avoid precipitation. Afterwards, the proteins were concentrated and further purified by Superose 6 Increase 10/300 GL (GE Healthcare) size exclusive chromatography.

#### **Verifications of the recombinant proteins by matrix assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF).**

The purified LC3B, HTTExon1Q25-MBP, and HTTExon1Q72-MBP proteins were dialyzed into 5 mM NH<sub>4</sub>Ac by Superose 6 Increase size exclusive chromatography for linear mode MALDI-TOF analysis on a Bruker FLEX MALDI-TOF instrument. A total of 1500-2500 scans were averaged for each spectrum using an accelerating voltage of 25 kV. Sinapinic acid (SA, Bruker, cat. no. 820135) was used as the matrices for protein and peptide analyses. SA was made into 20 mg/mL solutions in 70% ACN, 0.1% TFA. For the acquisition of spectra from 10,000 to 100,000 amu, 2  $\mu$ L of sample was mixed with 2  $\mu$ L of SA solution in an Eppendorf tube, and 2  $\mu$ L of the mixture was loaded onto the MALDI plate. The calibration peptides for this range were BSA (M + 66431) (Sigma, cat. no. A1933). All spectra were obtained in positive linear mode. The amount of full-length HTT proteins were limited, and thus not validated by MALDI-TOF. Instead, they were further purified by ion exchange and gel filtration chromatography, and validated by Coomassie-blue staining (Extended data Fig. 1e) and Western-blot (Fig. 4b).

#### **Verifications of the recombinant LC3B by X ray diffraction crystallography**

Since the deletion of G120 (lipidation site) stabilizes LC3B protein, we utilized LC3B $\Delta$ G120 protein to get high resolution diffraction data. Purified LC3B $\Delta$ G120 protein was concentrated in the following buffer: 20 mM HEPES pH7.5, 150 mM NaCl. The LC3B $\Delta$ G120 crystal was grown in reservoir solutions consisting of 0.16 M ammonium sulfate, 0.08 M sodium acetate pH 4.6, 20% (w/v) PEG4000, 20% (v/v) glycerol and 0.01 M Taurine.

#### **Refinement**

The X-ray diffraction data were collected at 100 K in the beamline BL17U1 and BL19U1, SSRF. The wavelength for data collection was 0.97892 Å. Diffraction images were indexed and processed by HKL2000. The structure of LC3B $\Delta$ G120 (PDB ID 6J04, 1.90Å) was solved by molecular replacement with the Phaser 2.8 program from the CCP4 crystallography package using the (PDB ID code 1UGM) as the search model. The refinement was performed by Refmac 5.5 and Phenix 1.14. There are no

668 Ramachandran outliers to report. The related figure was drawn using PyMOL 2.2.

669 **Compound-protein interaction measurements by oblique-incidence reflectivity**  
670 **difference (OI-RD)**

671 For high-throughput preliminary screening of target proteins, a SMM was  
672 assembled into a fluidic cartridge and washed *in situ* with a flow of 1× PBS to remove  
673 excess unbound small molecules. After washing, the SMM was scanned with a  
674 label-free OI-RD scanning microscope to image small molecules immobilized on glass  
675 slides. After it was blocked with 7,600 nM BSA in 1× PBS for 30 min, SMM was  
676 incubated with the target protein for 2 h. HTTexon1-Q25-MBP at a concentration of  
677 454 nM, HTTexon1-Q72-MBP at a concentration of 238 nM, and LC3B at a  
678 concentration of 680 nM were screened on separate fresh SMMs. OI-RD images were  
679 scanned for each operation, including washing, blocking, and incubation. The OI-RD  
680 difference images (images after incubation – images before incubation) were utilized  
681 for analysis, and vertical bright doublet spots indicated compounds that bind with  
682 target proteins in both replicates. 8F20 and 10O5 were identified to bind to  
683 HTTexon1-Q72-MBP and LC3B, but not HTTexon1-Q25-MBP. The binding was  
684 further confirmed by the kinetics measurements (see below).

685 To measure binding kinetics of target proteins with compounds, we prepared new  
686 SMMs consisting of 8F20, 10O5, and AN2. Six identical microarrays were printed on  
687 one glass slide and each compound was printed in triplicates in a single microarray.  
688 The printed small SMMs were assembled into a fluidic cartridge with each microarray  
689 housed in a separate chamber. Before the binding reaction, the slide was washed *in*  
690 *situ* with a flow of 1× PBS to remove excess unbound samples, followed by blocking  
691 with 7,600 nM BSA in 1× PBS for 30 min. For binding kinetics measurement, 1 × PBS  
692 was first flowed through a reaction chamber at a flow rate of 0.01 mL/min for 5 min to  
693 acquire the baseline. 1 × PBS was then quickly replaced with the probe solution of the  
694 target protein at a flow rate of 2 mL/min for 9 sec followed by a reduced flow rate at  
695 0.01 mL/min to have the microarray incubated in the probe solution under the flow  
696 condition for 35 min (association phase of the reaction). The probe solution was then  
697 quickly replaced with 1 × PBS at a flow rate of 2 mL/min for 9 sec followed by a  
698 reduced flow rate of 0.01 mL/min to allow dissociation of probe for 30 min  
699 (dissociation phase of the reaction). By repeating the binding reactions of the target  
700 protein at three different concentrations on separate fresh microarrays, binding curves  
701 of compounds with the target protein at three concentrations were recorded with  
702 scanning OI-RD microscope. Reaction kinetic rate constants were extracted by fitting  
703 the binding curves globally using 1-to-1 Langmuir reaction mode.

704 **Compound-protein interaction measurements by microscale thermophoresis**

705 **(MST)**

706 The purified recombinant proteins were dialyzed into 1× PBS, and then labelled  
707 according to the protocol of Protein labeling kit RED-NHS (Nanotemper, cat. no. L001).  
708 All the tested stock compounds (10 mM) dissolved in DMSO were also diluted into the  
709 same buffer for the final MST assay. The MST experiment was performed using  
710 Monolith NT.115 instrument (NanoTemper Technologies). 500 nM of labelled proteins  
711 were mixed with the indicated concentrations of candidate compounds in the reaction  
712 buffer containing 20 mM HEPES, pH 7.4, 150 mM NaCl. The MST data were then  
713 collected under 40% infrared laser power and 20% light-emitting diode power. The  
714 data were analyzed by Nanotemper analysis software (1.5.41) and the  $K_d$  was  
715 determined.

716 **cDNA plasmids for transfection in mammalian cells**

717 The pEX-GFP-hLC3WT plasmid was obtained from Addgene (#24987) to  
718 express LC3B. The pTT-HTTexon1-Q72-MBP-His and pTT-HTTexon1-Q25-MBP-His  
719 were generated by subcloning HTTexon1 cDNAs into the mammalian expression  
720 vector pTT-MBP-His and then transiently transfected into HeLa cells to express  
721 HTTexon1 proteins for the colocalization experiments. The polyQ-GFP sequences  
722 (expressing Met-polyQ-sfGFP) were *de novo* synthesized and subcloned into the  
723 pcDNA vector. All plasmids were sequence validated. For transient transfections, the  
724 cells were plated at 50% confluence. After 24 h, the cDNAs were transfected with  
725 Lipofectamine 2000 (ThermoFisher Scientific, cat. no. 11668019) using the forward  
726 transfection protocol provided by the manufacturer.

727 **Cell culture**

728 For mouse primary cortical neuron cultures, cortices were isolated from P0 pups  
729 following genotyping. Cortices were dissected into cold  $Ca^{2+}$ - and  $Mg^{2+}$ -free PBS  
730 buffer. Chopped small pieces were digested in solution containing 2.5% trypsin  
731 (Sigma, cat. no. P1005) and DNase I (0.1 mg/mL, Sigma, cat. no. D5025), for 20–30  
732 min at 37 °C. Tissues were transferred to 10% FBS containing DMEM (ThermoFisher  
733 Scientific, cat. no. 11965) to cease digestion. Neurons were then dissociated by  
734 trituration with fire-polished glass pipettes, collected by spinning and plated onto  
735 polylysine-coated dishes at  $4 \times 10^5$  cell/35 mm dish. The growth medium was  
736 composed of Neurobasal A medium (ThermoFisher Scientific, cat. no. 10888022) with  
737 1× B-27 (ThermoFisher Scientific, cat. no. 17504044) and 1× N2 supplement  
738 (ThermoFisher Scientific, cat. no. 17504048). Cytosine-arabinofuranoside (Sigma,  
739 cat. no. C1768) was added at 6  $\mu$ M to inhibit glial growth.

740 Some of the primary patient fibroblasts were obtained from HD patients (Q47,  
741 Q49, Q55) and healthy sibling (WT, Q19) controls in a Mongolian Huntington's

disease family. The HD Q68 fibroblast line was obtained from Coriell Cell Repositories (Camden, NJ, USA). The PD line was obtained from an idiopathic Parkinson's disease patient, and the SCA3 line was obtained from a SCA3 patient with the ATXN3 expansion mutation (Q74). The studies were approved by The Ethic Community of Institutes of Biomedical Sciences at Fudan University (#28) for obtaining the HD and wild-type patient fibroblasts, and by Huashan Hospital Institutional Review Board at Fudan University (#174) for obtaining the PD and SCA3 patient fibroblasts. Verbal and written consent was obtained from patients. The procedures were in compliance with all relevant ethical regulations. The immortalized fibroblasts were generated by infection of lentivirus expressing SV40T. For generation of iPSCs (iPSCs), the primary fibroblasts were transduced with the retroviral STEMCCA polycistronic reprogramming system (Millipore, cat. no. SCR548). The iPSCs were confirmed positive for Tra-1-81, Tra-1-60, SSEA-4 and Nanog by immunofluorescence and flow-cytometry. All four vector-encoded transgenes were found to be silenced and the karyotype was normal. iPSCs were cultured in E8 medium (ThermoFisher Scientific, cat. no. A1517001) on Matrigel (Corning, cat. no. 354277) surface. iPSCs were differentiated to Pax6-expressing primitive neuroepithelia (NE) for 10-12 days in a neural induction medium. Sonic hedgehog (SHH, 200 ng/ml) was added at days 10-25 to induce ventral progenitors. For neuronal differentiation, neural progenitor clusters were dissociated and placed onto poly-ornithine/laminin-coated coverslips at day 26 in Neurobasal medium (ThermoFisher Scientific, cat. no. 21103049), with 1× B-27 (ThermoFisher Scientific, cat. no. 17504044), 1× N-2 (ThermoFisher Scientific, cat. no. 17504048), brain derived neurotrophic factor (BDNF, 20 ng/ml, Protech, cat. no. 450-02), glial-derived neurotrophic factor (GDNF, 10 ng/ml, Protech, cat. no. 450-10), insulin-like growth factor 1 (IGF1, 10 ng/ml, Protech, cat. no. 100-11) and Vitamin C (Sigma cat. no. D-0260, 200 ng/ml). The mouse striatal cells (STHdh) were obtained from Coriell Cell Repositories (Camden, NJ, USA). The HEK293T cells and the HeLa cells were originally obtained from American Type Culture Collection (ATCC). STHdh, HeLa and HEK293T cells were cultured in DMEM (ThermoFisher Scientific, cat. no. 11965) with 10% (vol/vol) FBS (ThermoFisher Scientific, cat. no. 10082-147). Atg5 WT and KO MEFs were from N. Mizushima. All the mammalian cell lines were maintained at 37 °C incubator with 5% CO<sub>2</sub>, except STHdh cells, which were maintained at 33 °C with 5% CO<sub>2</sub>. The cells were tested every two months by a TransDetect PCR Mycoplasma Detection Kit (Transgen Biotech, cat. no. FM311-01) to ensure that they are mycoplasma free. The CellTiter-glo assay was performed to measure cell viability with the indicated compound treatment (Extended Data Fig. 2e) following the protocol provided in the kit (Promega, cat. no. G7570).

779 **HD *Drosophila* models**

780 The nervous system driver line *elav-GAL4* (*c155*), and the HTT-expressing lines  
781 *UAS-flHTT-Q16* and the *UAS-flHTT-Q128* (expressing human full-length HTT with  
782 16Q and 128Q, respectively, when crossed to the GAL4 line) lines were obtained from  
783 the Bloomington *Drosophila* Stock Center at University of Indiana  
784 (<http://flystocks.bio.indiana.edu/>), and maintained in a 25 °C incubator. Crosses were  
785 set up between virgin female flies carrying *elav-GAL4* driver and the *UAS-flHTT-Q16*  
786 or *UAS-flHTT-Q128* male flies to generate the desired genotypes.

787 **HD mouse models**

788 The generation and characterization of the Hdh140Q knock-in mice have been  
789 previously described<sup>19</sup>. Mice were group-housed (up to 5 adult mice *per* cage) in  
790 individually vented cages with a 12 h light/dark cycle. The mouse experiments were  
791 carried out following the ARRIVE (Animal Research: Reporting of *In Vivo* Experiments)  
792 guidelines, and they were in compliance with all relevant ethical regulations. The  
793 Animal Care and Use Committee of the School of Medicine at Fudan University  
794 approved the protocol used in animal experiments (Approval #20140904 and  
795 #20170223-005).

796 **Compound treatment in cells and animals**

797 The compounds utilized in this study were all commercially available, and quality  
798 controlled by the vendors using NMR. 10O5: GW5074 (DC Chemicals; cat.no.  
799 DC8810); 8F20: ispinesib (Selleck; cat.no. S1452); AN1:  
800 5-bromo-3-[(4-hydroxyphenyl)methylidene]-2,3-dihydro-1H-indol-2-one (Specs; cat.  
801 no. AN-655/15003575); AN2: 5,7-Dihydroxy-4-phenylcoumarin (ChemDiv; cat.no.  
802 D715-2435); GSK923295 (Selleck, cat.no. S7090), BAY1217389 (Selleck, cat.no.  
803 S8215), PLX-4720 (Selleck, cat.no. S1152), Dabrafenib (Selleck, cat.no. S2807),  
804 Sorafenib Tosylate (Selleck, cat.no. S1040), rapamycin (Sigma-Aldrich, cat. no.  
805 R8781).

806 For compound treatment in the cells, the compounds were diluted in culture  
807 medium to 10X concentrations and added to the plated cells: for primary cultured  
808 neurons and iPSC-derived neurons, the compounds were added 5 days after plating;  
809 for patient fibroblasts and other cell lines, the compounds were added 1 day after  
810 plating. The cells were then collected 2 days later for measurement of HTT levels. For  
811 detection of HTT-LC3 colocalization, the cells were fixed 4 h after compound  
812 treatment. For caspase-3 activation detection, the cells were stressed (BDNF removal  
813 for iPSC-derived neurons) 1 day after compound treatment, and tested at the  
814 indicated time points.



815 For compound treatment in the *Drosophila*, flies were maintained in standard  
816 maize food at 25 °C. For drug feeding, maize media was heated to 45 °C until liquid  
817 and distributed into vials. Compounds were freshly prepared in DMSO and added to  
818 the media. New adult flies were transferred to vials with 400 µL the control (DMSO) or  
819 compound-containing food, which was changed every other day.

820 For compound treatment in mice using intracerebral ventricular (icv) injection, the  
821 3-month-old mice were anesthetized using a small animal anesthesia machine  
822 (MSS-3, MSS International, Keighley, UK) by Isoflurane (1.5% solution). We surgically  
823 implanted each mouse with a guide cannula directed toward the lateral ventricle. The  
824 coordinates for implantation were determined utilizing “The Mouse Brain in  
825 Stereotaxic Coordinates” and the guide cannulas were placed at 0.6 mm posterior,  
826 1.5 mm lateral (left), and 1.7 mm dorsal with respect to bregma. A cap with stylus was  
827 then inserted into the guide cannula to seal its opening. Mice were then allowed to  
828 recover from surgery for a week before being treated. For injection, we first inserted  
829 an internal injector cannula so that it extended 0.5 mm beyond the tip of the guide  
830 cannula to reach the lateral ventricle. We then injected the mice through the internal  
831 injector cannula using a 25 µL syringe (Hamilton 1700 Series Microliter Syringes ,  
832 Bonaduz, GR, CH) at a flow rate of 0.25 µL/min powered by a syringe pump (KDS  
833 Legato 130, Holliston, MA, USA) to administer 2 µL of compounds-containing artificial  
834 cerebrospinal fluid (ACSF: 1 mM glucose, 119 mM NaCl, 2.5 mM KCl, 1.3 mM MgSO<sub>4</sub>,  
835 2.5 mM CaCl<sub>2</sub>, 26.2 mM NaHCO<sub>3</sub>, 1 mM NaH<sub>2</sub>PO<sub>4</sub>) at a concentration of 25 µM  
836 (containing 0.125% vol/vol DMSO). 2 µL ACSF containing equivalent amount of  
837 DMSO (0.125% vol/vol) was used as the control. The injector cannula was left in  
838 place for approximately 60 s to allow for diffusion before placing the caps with stylus  
839 back in guide cannulas.

840 For compound treatment in mice using intraperitoneal (ip) injection, each mouse  
841 was weighed. The compounds were diluted with 0.9% NaCl intravenous infusion  
842 solution to 0.05 µg/µL (containing 0.011 µg/µL DMSO) and injected into each mouse  
843 based on the weight of the mouse (500 µg/kg, containing 110 µg/kg DMSO). As  
844 controls, equivalent amount of DMSO was diluted and injected in the same way.  
845 Injection of 0.9% NaCl intravenous infusion solution alone was also tested and  
846 showed no difference (Extended data Fig. 9f-h). One injection *per* day was performed  
847 for two weeks before subsequent behavioral experiments or tissue extractions.

848 Note that in some of the experiments (Fig. 4-5 & Extended Data Fig. 6b-d), 8F20  
849 and/or AN1 were not tested. 8F20 was not tested because it did not have an effect *in*  
850 *vivo* by icv-injection (Extended Data Fig. 6a). AN1 was not tested because its

851 structure is highly similar as 10O5 while it had a weaker HTT-lowering effect by  
852 icv-injection (Extended Data Fig. 6a).

### 853 **Protein extraction from cells and tissues**

854 For protein extraction from cells, the cell pellets were collected and lysed on ice  
855 for 30 min in 1× PBS+1% Triton X-100+1× complete protease inhibitor (Sigma-Aldrich,  
856 cat. no. 11697498001), sonicated for 10 sec, and spun at >20,000 g at 4 °C for 15 min.  
857 The supernatants were then loaded and transferred onto nitrocellulose membranes  
858 for Western-blots. For mouse brain tissues, the mouse striata and cortices were  
859 dissected on ice and grinded by a tissue grinder for 5 min 60 Hz and lysed on ice for  
860 60 min in brain lysis buffer (50 mM Tris, 250 mM NaCl, 5 mM EDTA, 1% Triton X-100  
861 PH7.4) + 1× complete protease inhibitor (Roche, cat. no. 4693159001). The samples  
862 were then sonicated for 10 cycles, 15 s on and 20 s off, and then collected for  
863 Western-blots.

864 For protein extraction from the mouse brain, the brains were collected and the  
865 cortices were acutely dissected on ice and homogenized with a tissue grinder for 5  
866 min at 60 Hz and lysed on ice for 60 min in brain lysis buffer (50 mM Tris, 250 mM  
867 NaCl, 5 mM EDTA, 1% (vol/vol) Triton X-100, 1× complete protease inhibitor (Roche,  
868 cat. no. 4693159001), pH=7.4). The samples were then sonicated for 10 cycles, 15 s  
869 on and 20 s off, and then collected for Western-blots, HTRF or dot blots.

870 For mHTT measurements in the HD *Drosophila* model, the fly heads were  
871 collected at the age of 7 days and lysed on ice for 30 min in PBS + 1% (vol/vol) Triton  
872 X-100 + 1× complete protease inhibitor (Roche, cat. no. 4693159001), sonicated for  
873 10 cycles, 15 s on and 20 s off, and then collected for HTRF.

874 For all the samples, the protein concentrations were measured to correct the  
875 loadings. Different protein concentrations or cell numbers per well were tested to  
876 ensure that the signals were in the linear range. Background corrections were  
877 performed by subtracting the background signals from blank samples.

### 878 **Western-blot and filter trap assays**

879 For Western-blots, the samples were loaded onto the SDS page gel (5-12%  
880 depending on the molecular weight of the protein of interest). The proteins on the gel  
881 were then transferred to the nitrocellulose membranes for blocking and antibody  
882 detection. The signal was detected with ECL (Bio-Rad, cat. no. 1705061) after 1 h  
883 incubation of the membrane with secondary antibody 1:10,000.

884 The filter trap assay was performed similarly as previously described<sup>23</sup>, 2 µL (10  
885 µg) aliquots of each sample were loaded onto nitrocellulose membranes stacked in  
886 the Bio-Dot microfiltration apparatus (Bio-Rad). The membrane was blocked for 1 h  
887 with 5% milk and incubated overnight with the antibody 4C9 at a concentration of 1.5

888  $\mu\text{g}/\mu\text{l}$  in 5% milk diluted in PBS + 0.1% Tween-20. The signal was detected with ECL  
889 (Bio-Rad, cat. no. 1705061) after 1 h incubation of the membrane with secondary  
890 antibody 1:10,000.

#### 891 **Homogeneous Time Resolved Fluorescence (HTRF) assays**

892 For HTRF, the assays were similar as previously described<sup>25</sup>. The cell or tissue  
893 lysates were diluted with the original lysis buffer PBS + 1% (vol/vol) Triton X-100 + 1×  
894 complete protease inhibitor (Roche), utilized for lysing the samples, and then detected  
895 with indicated antibody pairs diluted in the HTRF assay buffer (50 mM  $\text{NaH}_2\text{PO}_4$ , 400  
896 mM NaF, 0.1% BSA, 0.05% (vol/vol) Tween-20, 1% (vol/vol) Triton X-100, pH 7.4).  
897 The donor antibody concentration was 0.023 ng/ $\mu\text{L}$  and the acceptor antibody  
898 concentration was 1.4 ng/ $\mu\text{L}$ , both in HTRF assay buffer. Different antibody pairs were  
899 used for different experiments as indicated in the figure legends. For all the samples,  
900 the signals were normalized to the total protein concentrations to ensure equal  
901 loadings. Different protein concentrations were pre-tested to ensure that the signals  
902 were in the linear range. Background corrections were performed by subtracting the  
903 background signals from blank samples.

#### 904 ***In vitro* c-Raf kinase assay**

905 *In vitro* c-Raf kinase assays were carried out with a c-Raf kinase assay kit (BPS  
906 Bioscience, cat. no. 79570). The assays were performed in a 96-well plate according  
907 to the manufacturing instruction. The samples and non-reactive negative controls  
908 were tested in duplicate according to the instruction.  
909 For details, 25  $\mu\text{L}$  of the mixture containing 5x kinase assay buffer (6  $\mu\text{L}$ ), ATP (1  $\mu\text{L}$ ),  
910 5x Raf substrate (10  $\mu\text{L}$ ) and water (8  $\mu\text{L}$ ) was added to a well. 5  $\mu\text{L}$  of water solution  
911 containing a test compound at a 10x desired concentration (DMSO was at 10% at the  
912 water solution) was added to the 25  $\mu\text{L}$  of mixture, and 20  $\mu\text{L}$  of 1x kinase assay  
913 buffer containing 2 ng/ $\mu\text{L}$  c-Raf kinase was added to the mixture in a well to initiate the  
914 kinase reaction (at this stage compounds were at 1x desired concentration, and  
915 DMSO was at 1% concentration). For a non-reactive negative control, 20  $\mu\text{L}$  of 1x  
916 kinase assay buffer containing no c-Raf was added to the mixture instead. The plate  
917 was incubated at 30 °C for 45 min. After the 45-min reaction, 50  $\mu\text{L}$  of kinase  
918 Kinase-Glo Max reagent (Promega, cat. no. V6071) was added to each well, and the  
919 plate was incubated at room temperature for 15 min, at dark. The plate was read with  
920 a microplate reader (BMG Labtech) for luminescence reading. The luminescence  
921 reading value measures the levels of ATP remaining, which is inversely related to  
922 kinase activity. The non-reactive negative control read value, indicating the level of  
923 initial added ATP, subtracted the level of ATP remaining (the luminescence reading)  
924 for the value of consumed ATP in the reaction that represents a kinase activity.

## 925 ***In vitro* pull-down assays**

926 We performed *in vitro* pull-down assays to test the compounds' influence on the  
927 HTT-LC3 interactions. The purified HTTexon1 (with the indicated tags), full-length  
928 proteins and the control proteins were incubated with amylose resin (New England  
929 BioLabs, cat. no. E8021L) at 4 °C for 30min. Immobilized amylose resins were then  
930 washed three times with HBS (20 mM HEPES pH7.5, 150 mM NaCl, 0.05%  
931 Tween-20). The resulting amylose resins containing about 10 µg of MBP-fused  
932 proteins were incubated with the indicated compounds (1 µM for 10O5 and 100 nM for  
933 AN2) or the DMSO control at the same volume in 300 µl of HBS at 4 °C for 1 h using  
934 sample mixer. 40 µg of purified LC3B protein were then added and incubated at 4 °C  
935 for another 2 h using sample mixer. The resin-bound proteins were eluted with 40 µl  
936 maltose buffer (10 mM maltose, 20 mM HEPES, 150 mM NaCl, pH 7.5) and then  
937 added with 20 µl SDS-PAGE sample loading buffer. Samples were then analyzed by  
938 SDS-PAGE and Western-blots.

939 GST pulldown was performed as the same procedures described above, except  
940 that GST-fused LC3B was immobilized onto magnetic conjugated GST mouse mAb  
941 beads (Cell Signaling Technology, cat.no.11847S) and eluted with SDS-PAGE  
942 protein loading buffer by vortex according to the instruction manual.

## 943 **Imaging-based autophagy assays**

944 Analysis of GFP-LC3 puncta for measuring autophagosomes: HeLa cells stably  
945 expressing GFP-LC3 were generated by transfection of pEGFP C1-LC3, and positive  
946 clones were selected by 500 µg/ml G418. The cells were then treated with vehicle  
947 (DMSO, 0.1%), 10O5, or AN2 for the indicated concentration, chloroquine (CQ, 20  
948 µM) treatment was used as a control. After 24 hours, cells were fixed in 4%  
949 paraformaldehyde (PFA) for 10 min. Images were acquired with confocal microscopy  
950 (Leica SP8) by the observer blinded to the identity of the slides. The number and size  
951 of GFP vesicles per cell was determined by Image J software. Images were  
952 processed with the despeckle function to decrease the noise, and a threshold was set  
953 to highlight puncta. Cells were selected by the freehand drawing tool. The  
954 analyze-particle function was used for the sizes and numbers of GFP puncta.

955 The mRFP-GFP-LC3 assay: this assay allows us to monitor autophagosome  
956 synthesis and maturation/fusion by labelling autophagosomes (green and red) and  
957 autolysosomes (red), since the low lysosomal pH in autolysosomes quenches the  
958 GFP signals<sup>32</sup>. HeLa cells stably expressing mRFP-GFP-LC3<sup>32</sup> were treated with  
959 vehicle (DMSO, 0.1%), 10O5, or AN2 for the indicated concentration, bafilomycin-A1  
960 (bafA1, 10 nM) treatment was used as a control. After 24 hours, cells were fixed in 4%  
961 PFA for 10 min. Images were acquired with confocal microscopy (Leica SP8) by the

observer blinded to the identity of the slides. The green and red single channel images were analyzed by Image J to quantify green and red puncta in the same way as in the GFP-LC3 assay described above.

#### **Detection of long-lived proteins by click-chemistry**

As an indicator of autophagy activity, the degradation of long-lived proteins was measured similarly as previously reported<sup>33</sup>. Basically, the HeLa cells with 70~80% confluency in a 6-well plate were washed with warm PBS and cultured in Met-free DMEM (ThermoFisher Scientific, cat. no.21013) added with dialyzed FBS for 1 h to deplete intracellular free Met reserves. The Met analog L-AHA (50  $\mu$ M) was then added to label the proteins for 18 h. After labeling, the cells were washed with PBS and cultured in regular culture medium containing 10x L-Met (2 mM) for 2 h to chase out short-lived proteins. The cells were then treated with the compounds versus the DMSO controls for 6 h before cell lysis and protein extraction. For the starvation sample, the culture medium was replaced with EBSS (ThermoFisher Scientific, cat. no. 24010043) for 6 h. The protein lysates were then used for the click reaction by the Click-it reaction kit (Click Chemistry tools, cat. no. C1001) following manufacturer's instructions, and the remaining L-AHA containing long-lived proteins were then conjugated with biotin. These proteins were then analyzed by electrophoresis and detected by the HRP-conjugated streptavidin (Beyotime, cat. no. A0303).

#### **Immunofluorescence and caspase-3 imaging**

For immunofluorescence of cultured cells, cells were fixed in 4% PFA for 10 min after washing with 1 $\times$  PBS for three times, and then washing and permeabilized in 0.5% (vol/vol) TritonX-100 for 10 min. The cells were then blocked in blocking buffer (4% BSA + 0.1% (vol/vol) Triton X-100 in 1 $\times$  PBS) for 30 min and incubated overnight at 4  $^{\circ}$ C with primary antibodies, and then washed three times with blocking buffer and incubated with secondary antibody at room temperature for 1 h. Coverslips were then washed three times, stained with 0.5 mg/ml DAPI for 5 min at room temperature, and then mounted in vectashield mounting medium (Vector, cat.no. H-1002). Images were taken by Zeiss Axio Vert A1 confocal microscopes and analyzed blindly by ImageJ for co-localization and TUBB3 quantifications. For co-localization experiments of transfected HeLa cells (Fig. 4c), the GFP signals were used to detect GFP-LC3B, and anti-His was used to detect HTTExon1-MBP-His proteins. Empty vector transfected cells were imaged to ensure the specificity of the signals. The co-localization was analyzed by calculating the ratio between overlapping puncta and the HTT (red) puncta for each cell, and the puncta numbers were counted blindly. For co-localization experiments of STHdh<sup>Q111/Q111</sup> cells, the endogenous mHTT protein was stained with the HTT antibody (Millipore, cat. no. MAB2166), and autophagosomes were stained

999 with the LC3B antibody (ThermoFisher Scientific, cat. no. 700712), which  
1000 preferentially detects LC3-II<sup>38</sup>. The co-localization was analyzed by ImageJ to  
1001 calculate the ratio between overlapping pixels and the HTT (red) positive pixels,  
1002 because the signals of the endogenous proteins were more dispersed and could not  
1003 be counted accurately. For TUBB3, the total area of TUBB3 signals and the DAPI  
1004 counts were analyzed by ImageJ. The former is then divided by the latter to calculate  
1005 the averaged area of TUBB3 in each neuron as an index for neurodegeneration *in*  
1006 *vitro*.

1007 For caspase-3 activity measurements of the iPSC-derived neurons, the NucView  
1008 488 caspase-3 dye (Biotium, cat. no. 30029) was used for the caspase 3 activity  
1009 detection as an indicator for apoptosis. The images were then taken every 3 h using  
1010 the Incucyte technology (Essen Bioscience, IncuCyte FLR), which takes images of 4  
1011 different fields in each well inside the cell culture incubator. The quantification was  
1012 performed by the Incucyte 2011A software, which identified the green fluorescent  
1013 puncta and quantified the fluorescent object count per field. The 4 fields *per* well were  
1014 quantified and averaged, and 4 independent wells were used for statistical analysis.

#### 1015 **Antibodies**

1016 Antibodies used for Western-blots, HTRF and/or  
1017 immunofluorescence/immunohistochemistry are as follows: the HTT antibodies 2B7<sup>39</sup>,  
1018 ab1<sup>40</sup> and MW1<sup>41</sup> have been described previously; commercially purchased  
1019 antibodies include HTT antibody 2166 (Millipore, cat. no. MAB2166), anti-polyQ  
1020 antibody 3B5H10 (Sigma, cat. no. P1874), anti-HTT antibody (D7F7)XP (Cell  
1021 Signaling Technologies, cat. no. 5656s), anti- $\beta$ -tubulin (Abcam, cat. no. ab6046),  
1022 anti-TUBB3 (Biolegends (previously Covance), cat. no. 801202), anti-ATXN3  
1023 (Millipore, cat. no. MAB5360); anti-Gapdh (Proteintech, cat. no. 60004-1), anti-NBR1  
1024 (ThermoFisher Scientific, cat. no. PA5-54660), anti- $\beta$ -actin (Beyotime, cat. no.  
1025 AA128); anti-TBP (Abcam, cat. no. ab818); anti-P62 (ThermoFisher Scientific, cat. no.  
1026 PA5-27247); anti-spectrin (Millipore, cat. no. MAB1622); anti-Ncoa4 (Santa cruz,  
1027 cat.no. sc-373739); anti-GST (ProteinTech, cat. no. HRP-66001); anti-GFP (Cell  
1028 Signaling Technologies, cat. no. 2956); anti-MBP (ProteinTech, cat. no. 15089-1-AP);  
1029 anti-His (Beyotime, cat. no. AH367); anti-BUBR1 (BD Transduction, cat.no, 612503);  
1030 anti-phospho-p44/42 MAPK (ERK1/2) and anti-phospho-MEK1/2 in the  
1031 Phospho-Erk1/2 Pathway Sampler Kit (Cell Signaling Technology, cat.no. 9911);  
1032 anti-LC3B (ThermoFisher Scientific, cat.no. PA1-16930 (for Western-blot) & cat. no.  
1033 700712 (for immunofluorescence)). All the antibodies used for immunofluorescence in  
1034 this study have been validated by knock-down experiments. All the HTT, polyQ and  
1035 ATXN3 antibodies used for HTRF and/or Western-blots have been validated by

1036 knock-down experiments and by comparing the signals from different genotypes in  
1037 previous studies from us and others. All the other antibodies have been validated by  
1038 previous literature or the vendor.

#### 1039 **Compound detection *in vivo* in brain tissue from ip-injected mice**

1040 The experiments were performed by the SIM-Servier joint laboratory. The mice  
1041 ip-injected with DMSO or the indicated compounds were anesthetized by chloral  
1042 hydrate (200  $\mu$ L/kg of 10% stock) at indicated time points, and the heart blood was  
1043 collected by vacuum blood collection tubes. The heart blood samples were further  
1044 spun at 10,000 rpm for 5 min to generate the heart plasma. The mice were then  
1045 perfused with 1X PBS to remove the blood. The mice were then sacrificed and the  
1046 brain samples were dissected. 5 times of volume of methanol: acetonitrile (50: 50,  
1047 vol/vol) were added to each sample, which was then homogenized. Following  
1048 ultrasonic treatment for 15 min, the homogenates were centrifuged for 5 min, then 20  
1049  $\mu$ L supernatant liquid was mixed with 20  $\mu$ L water for 30s before injection. Linear  
1050 range of 10O5 was 10-30000 ng/mL, and the linear range of AN2 was 0.3-10000  
1051 ng/mL. The LC-MS/MS analyses were performed on an Acquity ultra performance  
1052 liquid chromatography (UPLC) system (Waters Corporation, Milford, MA, USA)  
1053 coupled to a Xevo TQ-S mass spectrometer (Waters Corporation, Milford, MA, USA).  
1054 Chromatographic separation was performed using an Acquity UPLC BEH C18 (1.7  
1055  $\mu$ m 2.1  $\times$  50 mm) column supplied by Waters at a flow of 0.5 mL/min. Gradient elution  
1056 was used with a mobile phase composed of solvent A (water containing 0.1% formic  
1057 acid and 5 mM  $\text{NH}_4\text{AC}$ ) and solvent B (acetonitrile: methanol (9:1, vol/vol) containing  
1058 0.1% formic acid).

#### 1059 **Proteomics analysis**

1060 Samples were analyzed on Orbitrap Fusion Lumos mass spectrometers (Thermo  
1061 Fisher Scientific, Rockford, IL, USA) coupled with an Easy-nLC 1000 nanoflow LC  
1062 system (Thermo Fisher Scientific). Dried peptide samples were re-dissolved in  
1063 Solvent A (0.1% formic acid in water) and loaded to a trap column (100  $\mu$ m $\times$ 2 cm;  
1064 particle size, 3  $\mu$ m; pore size, 120  $\text{\AA}$ ; SunChrom, USA) with a max pressure of 280 bar  
1065 using Solvent A, then separated on a 150  $\mu$ m $\times$ 15 cm silica microcolumn (particle size,  
1066 1.9  $\mu$ m; pore size, 120  $\text{\AA}$ ; SunChrom, USA) with a gradient of 5–35% mobile phase B  
1067 (acetonitrile and 0.1% formic acid) at a flow rate of 600 nL/min for 75min. The FAIMS  
1068 device was placed before the mass spectrometer. FAIMS separation was performed  
1069 with the following settings: inner electrode temperature = 100  $^{\circ}\text{C}$ , outer electrode  
1070 temperature = 100  $^{\circ}\text{C}$ , carrier gas flow= 4.6 L/min, Dispersion Voltage = -5000 V,  
1071 entrance plate voltage = 250 V. The FAIMS carrier gas is  $\text{N}_2$  only. The noted CVs were  
1072 applied to the FAIMS electrodes. Each of the selected CVs was applied to sequential

1073 survey scans and MS/MS cycles (1s); the MS/MS CV was always paired with the  
1074 appropriate CV from the corresponding survey scan. For detection with Fusion or  
1075 Fusion Lumos mass spectrometry, a precursor scan was carried out in the Orbitrap by  
1076 scanning  $m/z$  300–1400 with a resolution of 120,000. The most intense ions selected  
1077 under top-speed mode were isolated in Quadrupole with a 1.6 $m/z$  window and  
1078 fragmented by higher energy collisional dissociation (HCD) with normalized collision  
1079 energy of 30%, then measured in the linear ion trap using the rapid ion trap scan rate.  
1080 Automatic gain control targets were  $5 \times 10^5$  ions with a max injection time of 50 ms for  
1081 full scans and  $1 \times 10^4$  with 35 ms for MS/MS scans. Dynamic exclusion time was set at  
1082 18 s. Data were acquired using the Xcalibur software (Thermo Scientific).

1083 Rawfiles were searched against the human National Center for Biotechnology  
1084 Information (NCBI) Refseq protein database (updated on 04-07-2013, 32,015 entries)  
1085 by Mascot 2.3 (Matrix Science) implemented on Proteome Discoverer 2.2 (Thermo  
1086 Scientific). The mass tolerances were 20 ppm for precursor and 0.5 Da for product  
1087 ions for Fusion Lumos. Up to two missed cleavages were allowed. The search engine  
1088 set cysteine carbamidomethylation as a fixed modification and N-acetylation,  
1089 oxidation of methionine as variable modifications. Precursor ion score charges were  
1090 limited to +2, +3, and +4. The data were also searched against a decoy database so  
1091 that protein identifications were accepted at a false discovery rate of 1%. Label-free  
1092 protein quantifications were calculated using a label-free, intensity-based absolute  
1093 quantification (iBAQ) approach.

1094 Proteins with at least 2 unique peptides with 1% FDR at the peptide level and  
1095 Mascot ion score greater than 20 were selected for further analysis. The file used for  
1096 protein inference and protein FDR calculation was derived from Mascot search results,  
1097 and the peptide spectrum match (PSM) was filtered via Percolator and customized  
1098 parameters, and then the proteins were assembled. The protein FDR was calculated  
1099 depending on the ratio of NPD (the number of assembled proteins from decoy  
1100 database searches) and NPT (the number of assembled proteins from target  
1101 database searches). The FOT was used to represent the normalized abundance of a  
1102 particular protein across samples. FOT was defined as a protein's iBAQ divided by the  
1103 total iBAQ of all identified proteins within one sample. The FOT was multiplied by  $10^5$   
1104 for the ease of presentation. Only the proteins detection in all compared samples  
1105 were utilized for comparison.

#### 1106 **Behavioral and lifespan experiments in HD *Drosophila* models**

1107 For behavioral experiments, we placed 15 age-matched virgin female flies in an  
1108 empty vial and tapped them down. The percentage of flies that climbed past a  
1109 7-cm-high line after 15 s was recorded. The mean of five observations is plotted for



each vial on each day, and data from multiple vials containing different batches of flies were plotted and analyzed by two-way ANOVA tests. The flies were randomly placed into each tube. For lifespan measurements, we placed 75 age-matched virgin female flies in an empty plastic vial and recorded the survival situation for each vial on each day. For both behavioral and lifespan measurement experiments, the person who performed the experiments were blinded to the drugs fed until data analysis.

#### **Behavioral experiments in HD mouse models**

All the behavioral experiments were performed during the light phase and the experimenters were blinded to the compound treatment and the genotype of each mouse. Both males and females have been used. All the mice were kept in the behavioral test room in dim red light for 1 h before starting the experiments. For rotarod experiments, mice were pre-trained on 3 consecutive days on the rotarod rotating at 4 rpm for 2 min. Mice were then tested for five days at an accelerating speed ranging from 4 to 40 rpm within 2 min. Each performance was recorded as the time in seconds spent on the rotating rod until falling off or until the end of the task. Each test included three repetitions with an inter-trial interval of 60 min in order to reduce stress and fatigue, and the means from these three runs were analyzed for each mouse. The balance beam test was run using a 2 cm thick meter stick suspended from a platform on both sides by metal grips. The total length is 100 cm. There was a bright light at the starting point and a dark box with food at the endpoint. The total time for each mouse to walk through the beam was recorded. For gripping force measurements, mice were allowed to grip the metal grids of a grip meter (Ametek Chatillon) with their forelimbs, and they were gently pulled backwards by the tail until they could no longer hold the grids. The peak grip strength observed in 10 trials was recorded.

#### **Statistics**

To ensure to reach a statistical power > 0.8, power analyses were performed for each assay based on estimated values by PASS 16 (<https://www.ncss.com/software/pass/>) before experiments. Estimation was based on our previously published results on similar experiments and preliminary experiments. The effect size was also estimated by Cohen's d, two means divided by the standard deviation for the data. The power analysis suggested  $n \geq 3$  for mHTT level measurements and  $n \geq 5$  for behavioral experiments. In all the experiments we performed, we have used a larger n than these numbers in case the effect was smaller than preliminary results, and we also performed post-experiment power analyses to ensure that power  $\geq 0.8$  for all the significant differences. Statistical comparisons between two groups were conducted by the unpaired two-tailed t tests.

Statistical comparisons among multiple groups were conducted by one-way ANOVA tests and post-hoc tests for the indicated comparisons (Dunnett's tests for comparison with a single control, and Bonferroni's tests for comparisons among different groups). Statistical comparisons for series of data collected at different time points were conducted by two-way ANOVA tests. The similarity of variances between groups to be compared was tested when performing statistics in GraphPad Prism 7 and Microsoft Excel 2016. Normality of data sets was assumed for ANOVA and t tests, and was tested by Shapiro-Wilk tests. When the data were significantly different from normal distribution, nonparametric tests were used for statistical analysis. All statistical tests were unpaired and two-tailed.

1157

## 1158 **Reference for Methods**

1159

- 1160 38 Hancock, M. K., Hermanson, S. B. & Dolman, N. J. A quantitative TR-FRET  
1161 plate reader immunoassay for measuring autophagy. *Autophagy* **8**, 1227-1244,  
1162 doi:10.4161/auto.20441 (2012).
- 1163 39 Weiss, A. *et al.* Single-step detection of mutant huntingtin in animal and human  
1164 tissues: a bioassay for Huntington's disease. *Anal Biochem* **395**, 8-15,  
1165 doi:10.1016/j.ab.2009.08.001 (2009).
- 1166 40 Sapp, E. *et al.* Native mutant huntingtin in human brain: evidence for  
1167 prevalence of full-length monomer. *J Biol Chem* **287**, 13487-13499,  
1168 doi:10.1074/jbc.M111.286609 (2012).
- 1169 41 Ko, J., Ou, S. & Patterson, P. H. New anti-huntingtin monoclonal antibodies:  
1170 implications for huntingtin conformation and its binding proteins. *Brain Res Bull*  
1171 **56**, 319-329, doi:10.1016/S0361-9230(01)00599-8 (2001).

1172

## 1173 **Acknowledgements**

1174 The authors would like to thank Drs. Junmei Lu, Min Jiang, Linyun Liu and Qian Huang for  
1175 their technical support of mouse behavioral experiments. Dr. Yanhui Xu for technical  
1176 supports of the protein purification. Dr. Hexige Saiyin for the help with obtaining human  
1177 patient fibroblasts. The authors would also like to thank the following funding supports:  
1178 National Key Research and Development Program of China (2016YFC0905100), National  
1179 Natural Science Foundation of China (8192500069, 81870990, 31961130379, 91649105,  
1180 31470764, 91527305, and 61505032), Science and Technology Commission of Shanghai  
1181 Municipality (18410722100), Natural Science Foundation of Shanghai (19ZR1405200),  
1182 Shanghai Municipal Science and Technology Major Project (No.2018SHZDZX01) and  
1183 ZJLab, and Hereditary Disease Foundation.

1184

### 1185 **Author Contribution**

1186 B Lu perceived the idea, initiated the project, designed experiments, analysed data and  
1187 wrote the manuscript. Y Fei and C Zhu performed the OI-RD screening and  $K_{on}/K_{off}$   
1188 measurements with data analysis. Y Ding, Z Wang, J Li, C Gao performed protein  
1189 purification, *in vitro* pull-down and structural biology experiments with data analysis. With  
1190 the help from others for blinding, Z Li performed the HTT measurements in cells and in  
1191 mouse models, the mouse phenotype experiments, the autophagy-related mechanistic  
1192 experiments, the control protein measurements, and neurotoxicity measurements. C Wang  
1193 replicated Z Li's HTT measurement and autophagy-related mechanistic experiments, and  
1194 performed additional HTT measurement and phenotypic experiments in HD fly models,  
1195 patient iPSC-derived neurons and MEFs, as well as all the measurements of other polyQ  
1196 proteins and all the MST experiments. Y Dang provided the compound library for the  
1197 screen. T Sha and C Ding performed proteomics experiments and analysis. S Luo and Y  
1198 Yang performed the measurements of autophagy function. L Ma, Y Sun and J Wang  
1199 provided and characterized the patient cell lines. X Sun did the initial subcloning of  
1200 full-length HTT and found the explanation for the observed "hook effects". C Lu  
1201 (biostatistician) performed biostatistical analysis. M Difiglia and Y Mei helped designing  
1202 the experiments and interpreting the data.

1203

### 1204 **Author Information**

1205 Information regarding reprints and permissions should be addressed to B Lu  
1206 ([luboxun@fudan.edu.cn](mailto:luboxun@fudan.edu.cn)). Correspondence and requests for materials should be  
1207 addressed to B Lu ([luboxun@fudan.edu.cn](mailto:luboxun@fudan.edu.cn)), or Y Fei ([fyy@fudan.edu.cn](mailto:fyy@fudan.edu.cn)), or Y Ding  
1208 ([yuding@fudan.edu.cn](mailto:yuding@fudan.edu.cn)). B Lu, Y Fei, Y Ding and Y Dang has filed two patents together  
1209 based on this study to the State Intellectual Property Office of China (201910180674.7  
1210 and 201910180717.1).

1211

### 1212 **Data Availability Statement**

1213 The protein structure data has been uploaded to the PDB database with entry number  
1214 6J04. Source data for all figure plots are provided with the paper. The full gel blots and the  
1215 proteomics datasets have been provided in supplementary information. The data that  
1216 support the findings of this study are available from the corresponding authors upon  
1217 reasonable request.

1218

### 1219 **Extended data figure/table legends**

#### 1220 **Extended Data Fig. 1. Protein purifications.**

1221 a) SDS-PAGE and linear mode MALDI-TOF mass spectrometry analysis of the  
1222 expression and purification of recombinant LC3B protein. For SDS-PAGE, lane 1, the

whole cell lysate before induction; lane 2, the whole cell lysate after induction; lane 3, the supernatants of induced cells; lane 4, the flow through fraction of Ni-NTA chromatography; lane 5, the eluates of Ni-NTA chromatography (GST-His8-LC3); lane 6, the eluates of GST-His8 tag removed LC3B by TEV protease; lane 7, the eluates of size exclusive chromatography; lane 8, molecular weight marker. The obtained mass spectrometry m/z peak of recombinant LC3B is 14660.811, consistent with theoretical calculations

b) Structural alignment of purified recombinant LC3BΔG120 (PDB: 6J04, yellow) with published LC3B structure (PDB: 1UGM, cyan) by Pymol.

c-d) SDS-PAGE and linear mode MALDI-TOF mass spectrometry analysis of the HTTexon1 proteins. For HTTexon1-Q72-MBP SDS-PAGE, lane 1, the supernatants of induced cells; lane 2, the insoluble fraction of induced cells; lane 3, the flow through fraction of Ni-NTA chromatography; lane 4, the eluates of Ni-NTA chromatography; lane 5, the eluates of size exclusive chromatography; lane 6, molecular weight marker. For HTTexon1-Q25-MBP SDS-PAGE, lane 1, molecular weight marker; lane 2, the induced cell lysate; lane 3, the supernatant fraction of induced cells; lane 4, the flow through fraction of Ni-NTA chromatography; lane 5 and 6, the eluates of Ni-NTA chromatography; lane 7 and 8, the eluates of size exclusive chromatography. The obtained mass spectrometry m/z peak of HTTexon1-Q72-MBP and HTTexon1-Q25-MBP are 64225.946 and 58228.893, consistent with theoretical calculations.

e) *Left* and *middle* panels: size exclusive chromatography of the recombinant full-length HTT-Q73 (flHTT-Q73) and HTT-Q23 (flHTT-Q23) proteins using Superose 6 5/150 GL. The major peak fractions were then collected pooled together for the SDS-PAGE analysis (*right* panel).

f) SDS-PAGE analysis of purified MBP-His8 (MBP), sfGFP (GFP), and Rpn10 proteins.

**Extended Data Fig. 2. Negative controls for OI-RD measurements and validation of the compounds' interaction with HTT and LC3 by MST.**

a) Similar to Fig. 1c-e, but for negative control proteins MBP-His8 (MBP), superfolder GFP (sfGFP), and ubiquitinated-Rpn10 (Ub-Rpn10). Association-dissociation curves of surface immobilized compounds 8F20 and 10O5 with these proteins were measured by OI-RD, and no compound-protein interactions were detected. For all association-dissociation curves, vertical dash lines mark the starts of association and dissociation phases of the binding event.

b) Binding of 10O5 and 8F20 to full-length HTT-Q73 (flHTT-Q73, black dots) or LC3B (red dots) in standard treated capillaries measured by MST. The compound bound protein fractions (bound / total) were calculated by the MST signals ( $F_{\text{norm}}$ ) at each compound concentrations, as well as the bound ( $F_{\text{norm\_bound}}$ , set as 100%) and the unbound ( $F_{\text{norm\_unbound}}$ , set as 0%) MST signals:  $\text{bound / total} = (F_{\text{norm}} - F_{\text{norm\_unbound}}) / (F_{\text{norm\_bound}} - F_{\text{norm\_unbound}}) \times 100\%$ . The fitted curves and calculated  $K_d$  values by the Nanotemper analysis software (1.5.41) for flHTT-Q73 and LC3B were indicated in each panel. Consistent with the OI-RD measurements (Fig. 1e), no binding was observed for the flHTT-Q23 protein (blue dots). The MST experiments were repeated > 3 times and showed consistent results.

c) Similar to b), except testing the compounds indicated in the X-axis. MST measurements

of the binding of indicated compounds to full-length HTT-Q73 (flHTT-Q73), full-length HTT-Q23 (flHTT-Q23) and LC3B in standard treated capillaries. The proteins tested were indicated in the legends.

d) Similar to Fig. 1c-e, but plotting the association-dissociation curves of surface immobilized compound AN2 with full-length HTT-Q73 (Q73), or full-length HTT-Q23 (Q23), or LC3B, or the negative control proteins MBP-His8 (MBP), superfolder GFP (sfGFP), and Rpn10.

For all association-dissociation curves, vertical dash lines mark the starts of association and dissociation phases of the binding event. The red dash lines are global fits to a Langmuir reaction model with the global fitting parameters listed at the bottom of each plot. No binding signals were observed for full-length HTT-Q23 proteins, and thus the parameters were not presented.

e) Cell viability measurement of cultured HD neurons measured by the CellTiter-glo assay. No toxicity was observed within the concentration range presented in Fig. 2, although the compound 8F20 became toxic to the cells when the concentration reached 300 nM.

1282

**Extended Data Fig. 3. mHTT lowering effects by mHTT-linker compounds could be detected by multiple antibodies and were dependent on autophagy.**

a) Representative Western-blots (HTT detected by the 2166 antibody) and quantifications of compound-treated cultured cortical neurons from Hdh<sup>Q7/Q140</sup> HD knockin mice. The neurons were treated with the indicated compounds (100 nM for 10O5, 8F20 and AN1; 50 nM for AN2) with or without the autophagy inhibitor NH<sub>4</sub>Cl (*upper panels*) or chloroquine (*lower left*), or the autophagy activator rapamycin (*lower right*). Same amount of culture medium was added in the controls (*upper panels*). The statistical analysis was performed by one-way ANOVA with post-hoc Dunnett's tests, and the F/degree of freedom/post-hoc p values have been indicated in each bar plot.

b) Western-blots using indicated HTT/polyQ antibodies for samples from cultured cortical neurons treated with the indicated compounds: 10O5 (100 nM), 8F20 (100 nM), AN2 (50 nM). The HTT gel blots presented in Fig. 2d (*right panel*) were cropped from first four blots. The low molecular weight bands were run out in these blots so that the wtHTT and mHTT could be better separated. Note that the weak bands just above 250 kDa in the first two blots were leftover signals from the Spectrin blotting. The Spectrin signals were too strong to be stripped completely.

c) Western-blots using the antibody MW1 or 3B5H10 that detects mHTT specifically. The relatively low molecular weight proteins were kept. No increase of potential polyQ-containing mHTT N-terminal fragments was observed.

d) HD patient iPSC-derived striatal neurons (Q47) were treated with the indicated compounds (100 nM, with 0.1% DMSO) in presence of an additional 0.1% DMSO or 10 mM NH<sub>4</sub>Cl, and the mHTT levels were measured by HTRF using the 2B7/MW1 antibody pair. All signals were normalized to the averaged signals from the DMSO control group. The statistical analysis was performed by one-way ANOVA with post-hoc Dunnett's tests, and the F/degree of freedom/post-hoc p values have been indicated in each bar plot. "\*\*\*\*": p<0.0001. The post-hoc analysis was not performed if the ANOVA tests did not show significance (p>0.05).

e) The HD patient immortalized fibroblasts (Q47) were transfected with the non-targeting control siRNA (Neg siRNA) or the ATG5 siRNA (target sequence: GCCUGUAUGUACUGCUUUA; ATG5 mRNA was knocked down to  $17.7 \pm 3.0\%$ ,  $n=3$ , as tested by qPCR), and then after 24 hours treated with the indicated compounds (100 nM) for another 48 hours. mHTT levels were then measured by HTRF using the 2B7/MW1 antibody pair. All signals were normalized to the averaged signals from the DMSO control group. The statistical analysis was performed by one-way ANOVA with post-hoc Dunnett's tests, and the F/degree of freedom/post-hoc p values have been indicated in each bar plot. "\*\*\*\*":  $p < 0.0001$ . The post-hoc analysis was not performed if the ANOVA tests did not show significance. The Western-blot of LC3 confirmed the partial inhibition of autophagy in the ATG5 knockdown cells.

f) Similar to e), but in wild-type (WT) or Atg5 knockout (Atg5 KO) mouse embryonic fibroblast lines (MEF) transfected with full-length mHTT (flHTT-Q73). The Western-blot of LC3 confirmed the inhibition of autophagy in the Atg5 KO cells.

For all panels, n indicates the number of independently plated wells, and bars represent mean and s.e.m.. Full-blot of cropped gels are shown in Supplementary Fig. 1.

**Extended Data Fig. 4. Potential influence of c-Raf and KSP pathways by treatment of the mHTT-LC3 linker compounds.**

a) Representative results (from 3 biological repeats) of the *in vitro* c-Raf kinase assay (see "Methods") showing that only 10O5 inhibits c-Raf activity within the concentration range tested.

b) Representative Western-blot and quantifications of phospho-MEK/ERK as an indicator of Raf activities (*left*) and phospho-BubR1 as an indicator of KSP inhibition (*right*) in cultured cortical neurons treated with indicated compounds (100 nM for 10O5, 8F20, AN1, and 50 nM for AN2) or the DMSO control.

c) Similar to b), but in immortalized HD patient fibroblasts (Q47). Note that the phospho-BubR1 is essentially absent and too weak to quantify, indicating that KSP was not inhibited by any of the compounds at the concentration tested.

For all panels, error bars represent mean and s.e.m.. For b, all data were corrected by the loading control ( $\beta$ -Tubulin) and then normalized to the averaged signal of the DMSO control group. The statistical analysis was performed by one-way ANOVA and the F/degree of freedom/p values have been indicated in each bar plot. The n number indicates the number of independently plated and treated wells.

**Extended Data Fig. 5. mHTT-LC3 linker compounds lowered mHTT in transgenic HD flies.**

a) Overlay between LC3B and predicted Atg8 structure showing high structural similarities.

b) Transgenic flies expressing full-length HTT-Q128 driven by *elav*-GAL4 were fed with indicated compounds at 10  $\mu$ M for 6 days, and their heads were then extracted for protein lysates. mHTT were then measured by HTRF using the 2B7/MW1 antibody pair.

Each dot represents the HTRF signal from each individual sample extracted from 5 fly heads. All the data were normalized to the average of the DMSO-fed control samples. The statistical analysis was performed by one-way ANOVA and Dunnett's post hoc tests. F (4,

31) = 15.67; “\*\*\*\*”:  $p < 0.0001$ .  
c) 10O5 (*upper* panel) and AN2 (*lower* panel) concentrations in heart plasma and brain tissues were measured by mass-spectrum at indicated time points for compound-injected mice (0.5 mg/kg). In the brain tissue, the 10O5 concentrations were ~20 to ~200 nM, and the AN2 concentrations were ~20 to ~40 nM, close to the effective doses that were capable of lowering mHTT in cultured neurons. Data were plotted by mean and s.e.m..

**Extended Data Fig. 6. mHTT-LC3 linker compounds lowered mHTT *in vivo* in the mouse brains.**

a) Western-blot (4 mice (3 months old) for each group) and quantifications of mHTT and wtHTT in the cortices from Hdh<sup>Q7/Q140</sup> knockin mice icv-injected with the indicated compounds (2  $\mu$ L at 25  $\mu$ M for each mouse) for 10 days at one-dose *per* day. HTT were detected by Western-blot using the 2166 antibody, and the statistical analysis was performed by one-way ANOVA and post-hoc Dunnett's tests. The F/degree of freedom/post-hoc p values have been indicated under each bar plot.

b) Similar to a), except that the compounds were delivered by ip-injection (0.5 mg/kg) to Hdh<sup>Q7/Q140</sup> mice at 5 months-old for 14 days at one-dose *per* day.

c) Similar to b), but from striata of ip-injected mice. The mice were injected at 10 months-old for 14 days at one-dose *per* day.

d) *Left* panel: representative dot-blot results (from two technical replicates) of the protein lysates in b) using the 4C9 antibody, which preferentially detects mHTT aggregates<sup>23</sup>. *Middle* panel: quantification of the dot-blot based on the averaged signals from two technical replicates. *Right* panel: measurement of mHTT aggregates by the 4C9/4C9 HTRF assay<sup>23</sup>.

For all panels, n indicates the number of mice tested, and bars represent mean and s.e.m.. For quantification, two to three technical replicates were averaged for each mouse. The statistical analysis was performed by one-way ANOVA with post-hoc Dunnett's tests, and the F/degree of freedom/post-hoc p values have been indicated in each bar plot.

**Extended Data Fig. 7. mHTT-LC3 linker compounds did not influence autophagy.**

a) HeLa cells stably expressing GFP-LC3B were treated with 2  $\mu$ L vehicle (0.1% DMSO), 10O5, or AN2 for the indicated concentration for 24 hours; chloroquine (CQ, 20  $\mu$ M) treatment was used as a control. After 24 hours, cells were fixed, and images were acquired with confocal microscopy. The number and size of GFP vesicles *per* cell was determined by Image J software (n number indicated on top of each plot). For each treatment, over 20,000 puncta were quantified (~100 puncta *per* cell in 226 cells). *Scale* bar: 10  $\mu$ m.

b) Representative images and quantifications of the numbers of autophagosomes (green<sup>+</sup> puncta) and autolysosomes (red<sup>+</sup>;green<sup>-</sup> puncta) in HeLa cells stably expressing mRFP-GFP-LC3B. *Scale* bar: 10  $\mu$ m. Autophagosome numbers or sizes were not influenced by 10O5 and AN2 at indicated concentrations after 24 hours treatment (or 4 hours treatment, not shown). The autophagosome fusion was also unaffected as indicated by the autolysosome number. Note that the autophagosome/autolysosome numbers/sizes were based on image analysis of the puncta, some of which may reflect multiple vesicles.

Green vesicles are considered to be autophagosomes (GFP+ puncta) and red vesicles are considered to be both autophagosomes and autolysosomes. The number of autolysosomes (RFP+ GFP- puncta) was achieved by subtracting the number of green vesicles from that of the red vesicles. >10,000 puncta from 194 cells were analyzed.

c) Representative Western-blots and quantifications of HeLa cells stably expressing GFP-LC3B. The “free GFP” was generated by lysosomal cleavage, and thus the “free GFP”/“GFP-LC3B” ratio was used as an index for autophagy flux, which was unaffected by 1005 or AN2, but decreased by the autophagy flux inhibitor chloroquine (CQ).

d) Representative Western-blots and quantifications of the chase signal of long-lived proteins in HeLa cells as an indicator of autophagy flux (see “Methods”). Consistent with previous reports<sup>33</sup>, starvation reduced the long-live protein chase signal, whereas rapamycin treatment had a milder effect. The mHTT-LC3 linker compounds 1005 and AN2 showed no influence in this assay.

e) Representative Western-blots and quantifications of LC3 in cultured cortical neurons treated with the indicated compounds. Normalized LC3-II/LC3-I was used as the indicator of autophagy. In the right panels, the 1005 and AN2 concentrations were 100 nM and 50 nM, respectively.

f) The SQSTM1/p62 levels were tested by Western-blots for the cortical tissues from mice injected by the indicated compounds or the DMSO control. Bars indicate mean and s.e.m.. For all panels, the n numbers indicated in each bar indicate the cell numbers (a-b), the number of independently plated wells (c-e) or mouse numbers (f). Error bars indicate mean and s.e.m.. The statistical analysis was performed by one-way ANOVA with post-hoc Dunnett’s tests for a-e, and by two-tailed unpaired t tests for f. Note that the post-hoc tests were not performed if the ANOVA tests failed to show significance. “\*\*\*\*”:  $p < 0.0001$  (post-hoc).

**Extended Data Fig. 8. Investigation on the specificity of mHTT lowering effects of mHTT-LC3 linker compounds.**

a) Representative Western-blots and quantifications of cultured cortical neurons treated with the indicated compounds. None of the proteins tested had an obvious change (> 10%).

b) Volcano plots of the proteomics analysis of cortices from ip-injected HD mice (10 m, 4 mice *per* group, injected for 14 days). The concentration injected was 0.5 mg/kg with 110 µg/kg DMSO, and equal amount of vehicle containing DMSO was injected as the controls. Only those proteins detected in both groups of samples used for comparisons were calculated and plotted. The dots plotted for HTT were pointed by red arrows. See Supplementary Table 2 for complete datasets. The bar plots indicate the total HTT levels normalized to the DMSO control. The actual mHTT reduction is anticipated to be more, because the compounds reduced mHTT allele-selectively.

c) Similar to b), but in cultured cortical neurons (from p0 pups, 3 wells *per* group). See Supplementary Table 3 for complete datasets.

In all panels, the error bars represent mean and s.e.m..


**Extended Data Fig. 9. mHTT-LC3 linker compounds lowered the mutant ATXN3**




1443 **protein with polyQ expansion in an allele-selective manner.**  
1444 a) Representative Western-blots and quantifications of ATXN3 levels in a SCA3 patient  
1445 fibroblast line treated with the indicated compounds. The lowering of mutant (Q74) but not  
1446 wild-type (Q27) ATXN3 was observed by treatment of linker compounds tested.  
1447 b) Quantification of the GFP intensity as an indicator of polyQ-sfGFP (25Q-GFP, 38Q-GFP,  
1448 46Q-GFP, and 72Q-GFP) protein levels in transfected HEK293T cells treated with the  
1449 indicated compounds using Incucyte. The lowering of 72Q-GFP, 46Q-GFP and 38Q-GFP,  
1450 but not 25Q-GFP was observed.  
1451 For both a) and b), the compound concentrations were 100 nM for 10O5 and AN1, and 50  
1452 nM for AN2. Bar plots present mean and s.e.m., and n numbers indicate independently  
1453 plated wells.  
1454 c) SDS-PAGE analysis of polyQ-sfGFP proteins (25Q, 38Q, 46Q, 53Q and 72Q) purified  
1455 from HEK293T cells. The protein purification methods were similar to the ones for HTT  
1456 proteins.  
1457 d) Binding of 10O5, AN1 and AN2 to sfGFP (GFP) or different polyQ-sfGFP (25Q-GFP,  
1458 38Q-GFP, 72Q-GFP) proteins in standard treated capillaries measured by MST,  
1459 performed and analyzed similarly as in Extended Data Fig. 2b. All these compounds  
1460 interact with 38Q-GFP and 72Q-GFP, but not 25Q-GFP or GFP.  
1461 e) Association-dissociation curves of surface immobilized compounds 10O5, AN1 and  
1462 AN2 with polyQ-sfGFP (72Q, 53Q, 46Q, 38Q and 25Q) proteins. For all  
1463 association-dissociation curves, vertical dash lines mark the starts of association and  
1464 dissociation phases of the binding event. The red dash curves are fits to a Langmuir  
1465 reaction model with the fitting parameters listed at the bottom of each plot. No binding  
1466 signals were observed for 25Q-sfGFP (25Q).  
1467 f-h) Results of mouse behavioral test performed similarly as in Fig. 5d-f, except that the  
1468 mice were injected with saline (0.9% NaCl) with DMSO (110 µg/kg) or without DMSO. The  
1469 statistical analysis was performed by two-way ANOVA with post-hoc Bonferroni's tests,  
1470 and the F/p values and degrees of freedom were indicated in the table underneath each  
1471 plot.  
1472 In all panels, the error bars represent mean and s.e.m..

1473  
1474 **Extended Data Table 1. Summary of data on mHTT lowering or rescue of**  
1475 **HD-relevant phenotypes**

1476 A summary table showing the percentage lowering of the mHTT or HTT levels, as well as  
1477 the percentage rescue of HD-relevant phenotypes (normalized to the difference between  
1478 HD and WT) in different HD models assayed by different approaches under optimal  
1479 conditions. The corresponding data figures have been indicated in the *middle* column. The  
1480 percentage change/rescue were presented in the form of mean ± s.e.m..  
1481

**Figure 1**

**Figure 2**



**Figure 3**






Figure 4


a




b



c










d



**Figure 5****a**

Human iPSC-derived neurons (TUBB3/DAPI; -BDNF)

**b** Human iPSC-derived neurons (-BDNF at 0 h)**c** *Drosophila* models expressing human full-length HTT**d** Mouse rotarod tests**e** Mouse balance-beam tests**f** Mouse gripping force tests**g** mHTT-LC3 linker compounds